

Ministerie van Landbouw, Natuur en
Voedselkwaliteit

> Retouradres Postbus 20401 2500 EK DEN HAAG

De Voorzitter van de Tweede Kamer
der Staten-Generaal
Postbus 20018
2500 EA DEN HAAG

**Directie Voedselkwaliteit en
Diergezondheid**

Cluster Voedselveiligheid
Prins Clauslaan 8
2595 AJ DEN HAAG
Postbus 20401
2500 EK DEN HAAG
www.minlnv.nl

T 070-3786868
F 070-3786113

Onze referentie
VD/2009/1201

Bijlagen
3

Datum 17 augustus 2009
Betreft Vrijwaringen ggo's

Geachte Voorzitter,

In het Algemeen Overleg biotechnologie van 18 juni jl. heb ik naar aanleiding van een verzoek van het lid Waalkens toegezegd om de Kamer te informeren als er sprake is van nieuwe vrijwaringsaanvragen in de komende 12 maanden. Ook alle openbare informatie rond vrijwaringen wordt aan de Kamer aangeboden.

De EU-regelgeving voor markttoelating van ggo's geeft lidstaten de mogelijkheid om het gebruik en/of de verkoop van in de Europese Unie (EU) toegelaten ggo's tijdelijk op hun grondgebied te beperken of te verbieden. Deze mogelijkheid wordt aangeduid met de term "vrijwaring". Lidstaten kunnen vrijwaringen instellen als nieuwe gegevens die wijzen op gevaren voor mens, dier of milieu daartoe aanleiding geven. Conform de Europese regelgeving dient vervolgens de Europese Commissie (EC) te beoordelen of de genomen vrijwaringen gerechtvaardigd zijn. De EC vraagt daarbij aan de Europese Voedselveiligheid Autoriteit (EFSA) om te beoordelen of de nieuwe gegevens wijzen op nieuwe risico's.

Bij deze bied ik u drie wetenschappelijke opinies van de EFSA aan over vrijwaringen die Oostenrijk heeft ingesteld om de in de EU toegelaten ggo-mais Mon863, de ggo-raapzaadvariëteiten GT73, MS8, RF3 en MS8xRF3 van het Oostenrijks grondgebied te weren. De EFSA concludeert in de drie opinies dat Oostenrijk geen nieuwe wetenschappelijke gegevens heeft aangeleverd die wijzen op gevaren voor mens, dier of milieu.

De EC zal mede op basis van de EFSA-opinies beoordelen of de genoemde Oostenrijkse vrijwaringen gerechtvaardigd zijn. Ik zal u dit oordeel van de Commissie zenden, zodra het beschikbaar is.

DE MINISTER VAN LANDBOUW, NATUUR EN
VOEDSELKWALITEIT,

G. Verburg

Leven
van het land,
geven
om natuur.

SCIENTIFIC OPINION

Request from the European Commission related to the safeguard clause invoked by Austria on oilseed rape GT73 according to Article 23 of Directive 2001/18/EC¹

Scientific Opinion of the Panel on Genetically Modified Organisms

(Question No EFSA-Q-2008-315)

Adopted on 15 June 2009

PANEL MEMBERS*

Hans Christer Andersson, Salvatore Arpaia, Detlef Bartsch, Josep Casacuberta, Howard Davies, Patrick du Jardin, Lieve Herman, Niels Hendriksen, Sirpa Kärenlampi, Jozsef Kiss, Gijs Kleter, Ilona Kryspin-Sørensen, Harry Kuiper, Ingolf Nes, Nickolas Panopoulos, Joe Perry, Annette Pöting, Joachim Schiemann, Willem Seinen, Jeremy Sweet and Jean-Michel Wal.

SUMMARY

On 27 July 2007, Austria invoked Article 23 of Directive 2001/18/EC (safeguard clause) to provisionally prohibit the marketing of genetically modified oilseed rape GT73 on its territory. Austria provided detailed reasons listed in supporting documents.

On 17 April 2008, the European Food Safety Authority (EFSA) has been requested by the European Commission to provide a scientific opinion on the statement and documents submitted by Austria in the context of a safeguard clause invoked under Article 23 of Directive 2001/18/EC.

In light of the information package provided by Austria in support of its safeguard clause and, having considered all relevant scientific publications, the Scientific Panel on Genetically Modified Organisms (GMO Panel) of EFSA concludes that, in terms of risk to human and animal health and the environment, no new scientific evidence was presented that would invalidate the previous risk assessment of oilseed rape GT73. The EFSA GMO Panel also concludes that no new scientific data or information was provided in support of adverse effects of oilseed rape GT73 on the environment and on human and animal health in Austria.

¹ For citation purposes: Scientific Opinion of the Panel on Genetically Modified Organisms on a request from the European Commission related to the safeguard clause invoked by Austria on oilseed rape GT73 according to Article 23 of Directive 2001/18/EC. *The EFSA Journal* (2009) 1151, 1-16.

* (minority opinion) This opinion is not shared by 0 members of the Panel. / (conflict of interest) 0 members of the Panel did not participate in (part of) the discussion on the subject referred to above because of possible conflicts of interest.

Therefore, no specific scientific evidence, in terms of risk to human and animal health and the environment, were provided that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC.

Key words: GMOs, oilseed rape (*Brassica napus*), GT73, Austria, safeguard clause, human and animal health, environment, Directive 2001/18/EC

TABLE OF CONTENTS

Panel Members	1
Summary	1
Table of Contents	3
Background	4
Terms of reference as provided by the European Commission.....	4
Acknowledgements	4
Assessment	5
1. Introduction	5
2. Assessment of documents provided by Austria.....	5
2.1. Food and feed safety issues.....	6
2.1.1. Toxicological and allergenicity assessment.....	6
2.2. Environmental safety issues and post-market environmental monitoring	6
2.2.1. Environmental risk assessment.....	6
2.2.2. Post-market environmental monitoring plan	9
2.2.3. Conclusion.....	9
Overall Conclusions and Recommendations.....	9
Documentation provided to EFSA	10
References	10

BACKGROUND

On 27 July 2007, Austria notified to the European Commission a national safeguard clause on genetically modified (GM) oilseed rape event GT73 under Article 23 of Directive 2001/18/EC. The notification was accompanied by the scientific document entitled “*Scientific arguments for an import ban of herbicide tolerant oilseed rape GT73 (notification C/NL/98/11)*”.

On 17 April 2008, the European Food Safety Authority (EFSA) has received a request from the European Commission to provide a scientific opinion from its Scientific Panel on Genetically Modified Organisms (GMO Panel) on the statement and documents submitted by Austria in the context of its invoked safeguard clause.

TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

EFSA was requested, under Article 29(1) and in accordance with Article 22(5) of Regulation (EC) No 178/2002, to provide a scientific opinion as to “*whether, in accordance with Article 23 of Directive 2001/18/EC, the statement and documents submitted by the Austrian authorities comprise new or additional information affecting the environmental risk assessment, such that detailed grounds exist to consider that the above authorised GMO, for the uses laid down in the corresponding consent, constitute a risk to human health or the environment*”.

ACKNOWLEDGEMENTS

The European Food Safety Authority wishes to thank the members of the Working Groups on Molecular Characterisation, Food/Feed and Environment, as well as the following members of its staff: Yann Devos and Sylvie Mestdagh for the preparation of this opinion.

ASSESSMENT

1. Introduction

Directive 2001/18/EC provides the possibility for Member States to invoke safeguards on specific genetically modified organisms in the case where new or additional information, made available since the date of the consent, would affect the risk assessment of an authorised GMO. Provisions foreseen by Austria seek to provisionally prohibit the marketing of oilseed rape GT73 for its intended uses on the Austrian territory.

The EFSA GMO Panel examined the set of supporting documents submitted by Austria. In this respect, the EFSA GMO Panel assessed whether the submitted documents comprise new scientific information that would change the outcome of previously performed risk assessments, and if detailed grounds exist to consider that the authorised oilseed rape GT73, for its intended uses, constitute a risk to human and animal health or the environment.

The EFSA GMO Panel looked for evidence for GMO-specific risks – including long-term effects (e.g., BEETLE report, 2009) – taking into consideration the EFSA GMO Panel guidance document for the risk assessment of GM plants and derived food and feed (EFSA, 2006a) as well as any related risk assessments carried out in the past. In addition, the EFSA GMO Panel considered the relevance of raised concerns in light of the most recent scientific data and relevant peer-reviewed publications.

2. Assessment of documents provided by Austria

A set of supporting documents, accompanying the mandate of the European Commission (see *Terms of Reference as provided by the European Commission*), was forwarded to EFSA on 17 April 2008.

Austria provided the following set of documents in support of its safeguard clause:

- Austrian letter on new supplementary scientific evaluation (13 July 2007);
- Verbot des Inverkehrbringens von gentechnisch verändertem Raps aus der Öraplinie GT73 in Österreich;
- Pascher, K., Narendja, F., Rau, D. (2006). Feral oilseed rape – Investigations on its potential for hybridisation, Studie im Auftrag des Bundesministeriums für Gesundheit und Frauen, Forschungsberichte der Sektion IV, Band 3/2006;
- Scientific arguments for an import ban of herbicide tolerant oilseed rape GT73 (Notification C/NL/98/11).

Based on the supporting documents, several issues were identified and therefore considered by the EFSA GMO Panel in the following two main areas: (1) the toxicological and allergenicity risk assessment, and (2) the environmental risk assessment and post-market environmental monitoring plan relating to the accidental spillage of oilseed rape GT73 seeds.

Issues related to the coexistence of oilseed rape cropping systems and the adventitious presence of authorised GM material in non-GM products were not considered, as they fall

outside the remit of the EFSA GMO Panel. In addition, the EFSA GMO Panel notes that it only gives its opinion on the scientific quality of the post-market environmental monitoring activities proposed by applicants, whilst the final endorsement thereof is done by risk managers.

During its assessment, the EFSA GMO Panel identified issues raised by the Austrian authorities that would require further clarifications. To present and clarify the provided set of documents, an informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel and EFSA staff was held on 23 April 2009. A representative of the European Commission attended the meeting as observer.

2.1. Food and feed safety issues

2.1.1. Toxicological and allergenicity assessment

The EFSA GMO Panel observes that the two publications related to toxicological and allergenicity aspects of the risk assessment quoted by Austria (Spök et al., 2004, 2005) do not provide new data specific on the safety of oilseed rape GT73: current approaches for assessing toxicological and allergenicity risks of genetically modified organisms are questioned in generic terms. The EFSA GMO Panel emphasises that the approach taken by the Panel in order to assess the potential toxicological and/or allergenicity risks of GM plants is in accordance with internationally developed guidelines (e.g., Codex Alimentarius, 2003).

With regard to toxicological assessment, Austria refers to various arguments provided by the applicant in its technical report that have already been addressed in the previous EFSA GMO Panel opinion on oilseed rape GT73 (EFSA, 2004). Because the compositional analysis of oilseed rape GT73 gave no indications of any changes, no further *in vitro* or *in vivo* toxicological testing was deemed necessary.

The general perspectives on the allergenicity assessment of GM food and feed to which Austria refers are currently being considered by a dedicated working group of the EFSA GMO Panel. The approach followed by the EFSA GMO Panel in its assessment of oilseed rape GT73 and the transgenic proteins within the latter aligns with the 'weight of evidence' approach described in the EFSA GMO Panel guidance document for risk assessment of GM plants and derived food and feed products (EFSA, 2006a) and in the guidelines of Codex Alimentarius (Codex Alimentarius, 2003), which have also been endorsed by Austria.

The EFSA GMO Panel concludes that the toxicological and allergenicity information provided by Austria are not new and have already been considered in the respective opinions of the EFSA GMO Panel.

2.2. Environmental safety issues and post-market environmental monitoring

2.2.1. Environmental risk assessment

In line with its previous scientific opinions on herbicide tolerant oilseed rape GT73 (EFSA, 2004), MS8xRF3 (EFSA, 2005) and T45 (EFSA, 2008), the EFSA GMO Panel confirms that in regions where oilseed rape is grown and/or where oilseed rape seeds are imported and transported, feral oilseed rape populations are likely to occur in non-natural disturbed

ecosystems (such as ports, processing facilities, margins of agricultural fields, roadside verges, railway lines, and wastelands) (Bagavathiannan and Van Acker, 2008). It is well-known that human activity contributes to the dispersal of plants (Wichmann et al., 2009), especially the transport of seeds by vehicles (Zwaenepoel et al., 2006; von der Lippe and Kowarik, 2007a,b; Garnier et al., 2008).

In the scientific literature, the occurrence of feral oilseed rape populations has been reported not only in Austria (Pascher et al., 2006), but also in Canada (Simard et al., 2002; Yoshimura et al., 2006; Knispel et al., 2008), France (Pessel et al., 2001; Garnier et al., 2008; Pivard et al., 2008a,b), Germany (Menzel, 2006; Reuter et al., 2008; Neuffer, 2009), Japan (Saji et al., 2005; Aono et al., 2006; Kawata et al., 2008; Nishizawa et al., 2009) and the United Kingdom (Crawley and Brown, 1995, 2004; Wilkinson et al., 1995; Charters et al., 1999; Norris and Sweet, 2002). These populations can be large and show significant variation in size from one year to the next (Crawley and Brown, 1995).

Due to its early germination potential and capacity to capture resources rapidly, oilseed rape can take advantage of disturbed land (Blackshaw et al., 2003, 2004). However, successful recruitment of oilseed rape from seed spillage from passing traffic mainly depends upon its ability to compete for space with primary colonizers, particularly perennial grasses. In most non-agricultural areas, oilseed rape lacks the ability to establish stable populations due to the absence of competition-free gaps (Crawley et al., 1993; Warwick et al., 1999; Hails et al., 2006). Once established, oilseed rape populations often become extinct after 2 to 4 years (Crawley and Brown, 1995; Crawley et al., 2001; Norris and Sweet, 2002). If habitats are disturbed on a regular basis (e.g., by mowing, herbicide application, soil disturbance) and replenished with seed from seed spillage or recruitment from seeds produced by residents or from seeds from the seedbank, then feral oilseed rape populations can persist for longer periods (8-10 years) (Pessel et al., 2001; Pivard et al., 2008a,b). Using genetic analyses and farmer surveys, Pessel et al. (2001) revealed that some members of feral oilseed rape populations in road verges in France originated from varieties that had not been marketed for at least 8 years.

Oilseed rape is generally regarded as an opportunistic species, and not as an environmentally hazardous colonizing species (Warwick et al., 1999). Several field studies and model predictions reported that the presence of herbicide tolerance in oilseed rape does not confer a fitness advantage, unless the respective herbicide is applied (Crawley et al., 1993, 2001; Fredshavn et al., 1995; Warwick et al., 1999, 2004; Norris and Sweet, 2002; Claessen et al., 2005a,b; Simard et al., 2005; Garnier and Lecomte, 2006; Garnier et al., 2006). In the absence of glyphosate-containing herbicide applications, oilseed rape GT73 is neither more likely to survive, nor more invasive or persistent than its conventional counterpart. Moreover, there is no evidence that tolerance to glyphosate enhances seed dormancy, and hence the persistence of feral oilseed rape populations (Claessen et al., 2005a,b; Sweet et al., 2004; Lutman et al., 2005, 2008; Messéan et al., 2007). Only where and when glyphosate-containing herbicides are applied, is oilseed rape GT73 expected to have a fitness advantage. In this respect, the scientific information provided in the Austrian safeguard clause notification does not give any new information regarding increased likelihood of establishment or survival of feral oilseed rape plants in case of accidental release into the environment of oilseed rape GT73 seeds during transportation and processing.

The EFSA GMO Panel is aware that if feral oilseed rape plants derived from spilled seeds remain uncontrolled and reproduce, they may survive, outcross and eventually disperse genes

to cross-compatible plants such as *Brassica rapa* and *Raphanus raphanistrum* (Scheffler and Dale, 1994; Eastham and Sweet, 2002; Chèvre et al., 2004; Warwick et al., 2003, 2004, 2008; Claessen et al., 2005b; Jørgensen, 2007; BEETLE report, 2009; Devos et al., 2009; Jørgensen et al., 2009). Scientific evidence suggests that feral oilseed rape populations may serve as a reservoir that could hold and return (trans)genes to cultivated populations of oilseed rape in a different place and time, and act as a genetic bridge delivering the (trans)genes to sympatric cross-compatible plants (Saji et al., 2005; Aono et al., 2006; Pascher et al., 2006; Yoshimura et al., 2006; Knispel et al., 2008; Nishizawa et al., 2009). In Canadian regions where GM oilseed rape is frequently grown (e.g., Beckie et al., 2006), feral oilseed rape populations were shown to actively outcross with cultivated populations of GM oilseed rape and to accumulate transgenes (Knispel et al., 2008). However, compared to cultivated oilseed rape populations, the contribution of feral oilseed rape plants in vertical gene flow is expected to be limited: feral oilseed rape populations are small compared to cultivated populations and contribute little to the pollen load in the environment (Colbach et al., 2001a,b, 2005; Devaux et al., 2005, 2007, 2008; Gruber and Claupein, 2006; Knispel et al., 2008; Colbach, 2009). Recent observations in Canada confirmed the persistence of a herbicide tolerance trait over a period of 6 years into a population of *B. rapa* in the absence of herbicide selection pressure (with the exception of possible exposure to the active substance glyphosate in 1 year) and in spite of fitness costs associated with hybridisation (Warwick et al., 2008). However, so far, there are no compelling data to suggest that the presence of an herbicide tolerance trait in a wild relative changes the behaviour of the wild relative. In the absence of glyphosate-containing herbicides, hybrids or wild relatives containing the herbicide tolerance trait do not show any enhanced fitness and behave as conventional plants. Thus escaped plants and genes dispersed to other cross-compatible plants would not create additional environmental impacts. If needed, feral oilseed rape GT73 and hybridised/introgressed relatives can be managed by the use of other herbicides and/or adequate mechanical practices (Beckie et al., 2004; Devos et al., 2004; Warwick et al., 2004; Légère, 2005; Simard et al., 2005; Gruber et al., 2008; Lutman et al., 2008).

The environmental exposure due to GM oilseed rape grain imports is anticipated to be low, as the amounts of viable oilseed rape seeds imported in the EU are limited with most seeds being imported by boat and crushed in or near the ports of entry. Some of the oilseed rape seeds imported into the EU are likely to be transported inland to Austria by boat. Moreover, some oilseed rape seeds entering Austria are transported by road or rail to processing plants. As indicated above, survival and outcrossing from plants derived from seed spillage will be at very low frequencies and have no hazardous environmental consequences compared to current feral oilseed rape populations.

In conclusion, the EFSA GMO Panel confirms that feral oilseed rape plants are likely to occur wherever oilseed rape is cultivated and/or transported and that transgenic oilseed rape is no exception (e.g., Saji et al., 2005; Aono et al., 2006; Yoshimura et al., 2006; Bagavathiannan and Van Acker, 2008; Kawata et al., 2008; Knispel et al., 2008; Nishizawa et al., 2009). However, there is no evidence that the herbicide tolerance trait introduced by genetic engineering results in increased invasiveness of oilseed rape GT73, except when glyphosate-containing herbicides are applied. As such, escaped plants and genes dispersed to other cross-compatible plants would not create additional agronomic or environmental impacts. This – together with the assessment that oilseed rape GT73 and hybridising relatives have no enhanced fitness or invasiveness characteristics (except in the presence of glyphosate-containing herbicides) – confirms earlier conclusions of the EFSA GMO Panel.

2.2.2. Post-market environmental monitoring plan

The EFSA GMO Panel maintains its position that the scope of the post-market environmental monitoring plan provided by the applicant complies with (1) the intended uses of oilseed rape GT73, which excludes cultivation, (2) the requirements of the EFSA GMO Panel guidance document for the risk assessment of GM plants and derived food and feed products (EFSA, 2006a), and (3) the EFSA GMO Panel scientific opinion on post-market environmental monitoring (EFSA, 2006b).

European operators importing, handling and processing viable oilseed rape commodities have recently joined with the European Association of Bioindustries (EuropaBio) in developing monitoring systems for GM oilseed rape imported through main points of entry and processing facilities (Lecoq et al., 2007; Windels et al., 2008). These monitoring systems are based on Hazard Analysis and Critical Control Points (HACCP) principles. Therefore, it is anticipated that risk managers will opt for post-market environmental monitoring of imported oilseed rape GT73 seeds in accordance with these arrangements.

2.2.3. Conclusion

The EFSA GMO Panel confirms its opinion that the likelihood of unintended environmental effects as a consequence of spread of (trans)genes from oilseed rape GT73 will not differ from that of conventional oilseed rape varieties in the context of its intended uses. The EFSA GMO Panel agrees with the monitoring plan submitted by the applicant, especially now that comprehensive arrangements have been made by applicants and operators for monitoring at major points of import and processing in the EU. However, the EFSA GMO Panel continues to recommend that appropriate management systems should be in place to minimise accidental loss and spillage of transgenic oilseed rape seeds during transportation, storage and handling in the environment, and processing into derived products. These conclusions are in line with previous scientific opinions of the EFSA GMO Panel on the import and processing of herbicide tolerant oilseed rape GT73 (EFSA, 2004), MS8xRF3 (EFSA, 2005) and T45 (EFSA, 2008).

OVERALL CONCLUSIONS AND RECOMMENDATIONS

The EFSA GMO Panel has investigated the claims and documents submitted in support of the Austrian safeguard clause and presented at the informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel and EFSA staff on 23 April 2009. In these documents, the EFSA GMO Panel did not identify any new data subject to scientific scrutiny or scientific information that would change previous risk assessments conducted on oilseed rape GT73 which currently has marketing consent in the EU. In addition, the Austrian submission did not supply scientific evidence, that the environment or ecology of Austria was different from other regions of the EU, sufficient to merit separate risk assessments from those conducted for other regions in the EU.

Having considered the overall information package submitted by Austria as well as a broad range of relevant scientific literature, the EFSA GMO Panel is of the opinion that there is no specific scientific evidence, in terms of risk to human and animal health and the environment, that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC for the marketing of oilseed rape GT73 for its intended uses in Austria. In conclusion, the EFSA GMO Panel finds that the scientific evidence currently available does

not sustain the arguments provided by Austria, and therefore the EFSA GMO Panel reiterates its previous scientific opinion on oilseed rape GT73.

DOCUMENTATION PROVIDED TO EFSA

1. Letter, dated 17 April 2008, with supporting documents from M.P. Carl, Director-General Environment EC, to Catherine Geslain-Lanéelle, Executive Director EFSA (ref ENV/B3/YK/gm D(2008) 518) requesting for a scientific opinion on the safeguard notification submitted by Austria under Article 23 of Directive 2001/18/EC for oilseed rape GT73 and comprising the following supporting documents:
 - Austrian letter on new supplementary scientific evaluation (13 July 2007);
 - Verbot des Inverkehrbringens von gentechnisch verändertem Raps aus der Ölrapslinie GT73 in Österreich;
 - Pascher, K., Narendja, F., Rau, D. (2006). Feral oilseed rape – Investigations on its potential for hybridisation, Studie im Auftrag des Bundesministeriums für Gesundheit und Frauen, Forschungsberichte der Sektion IV, Band 3/2006;
 - Scientific arguments for an import ban of herbicide tolerant oilseed rape GT73 (Notification C/NL/98/11).
2. Letter, dated 15 October 2008, from Catherine Geslain-Lanéelle, Executive Director EFSA, to Jos Delbeke, Acting Director-General Environment EC (ref CGL/RM/PB/SM/shv (2008) 3369831), acknowledging the receipt of the mandate accompanied with the supporting documents.

REFERENCES

- Aono, M., Wakiyama, S., Nagatsu, M., Nakajima, N., Tamaoki, M., Kubo, A., Saji, H., 2006. Detection of feral transgenic oilseed rape with multiple-herbicide resistance in Japan. *Environmental Biosafety Research*, 5: 77-87.
- Bagavathiannan, M.V., Van Acker, R.C., 2008. Crop ferality: implications for novel trait confinement. *Agriculture, Ecosystems and Environment*, 127: 1-6.
- Beckie, H.J., Séguin-Swartz, G., Nair, H., Warwick, S.I., Johnson, E., 2004. Multiple herbicide-resistant canola (*Brassica napus*) can be controlled by alternative herbicides. *Weed Science*, 52: 152-157.
- Beckie, H.J., Harker, K.N., Hall, L.M., Warwick, S.I., Légère, A., Sikkema, P.H., Clayton, G.W., Thomas, A.G., Leeson, J.Y., Séguin-Swartz, G., Simard, M.-J., 2006. A decade of herbicide-resistant crops in Canada. *Canadian Journal of Plant Science*, 86: 1243-1264.
- BEETLE report, 2009. Long-term effects of genetically modified (GM) crops on health and the environment (including biodiversity): prioritization of potential risks and delimitation of uncertainties. German Federal Office of Consumer Protection and Food Safety, BLaU-Umweltstudien and Genius GmbH, http://ec.europa.eu/environment/biotechnology/pdf/beetle_report.pdf

- Blackshaw, R.E., Brandt, R.N., Janzen, H.H., Entz, T., Grant, C.A., Derksen, D.A., 2003. Differential response of weed species to added nitrogen. *Weed Science*, 51: 532-539.
- Blackshaw, R.E., Brandt, R.N., Janzen, H.H., Entz, T., 2004. Weed species response to phosphorus fertilization. *Weed Science*, 52: 406-412.
- Charters, Y.M., Robertson, A., Squire, G.R., 1999. Investigation of feral oilseed rape populations, genetically modified organisms research report (No. 12), Department of the Environment, Transport and the Regions, <http://www.defra.gov.uk/environment/gm/research/reports.htm>
- Chèvre, A.M., Ammitzbøll, H., Breckling, B., Dietz-Pfeilstetter, A., Eber, F., Fargue, A., Gomez-Campo, C., Jenczewski, E., Jørgensen, R., Lavigne, C., Meier, M., den Nijs, H., Pascher, K., Seguin-Swartz, G., Sweet, J., Stewart, N., Warwick, S., 2004. A review on interspecific gene flow from oilseed rape to wild relatives. In: den Nijs, H.C.M., Bartsch, D., Sweet, J. (Eds), *Introgression from Genetically Modified Plants into Wild Relatives* (pp 235-251), CABI publishing.
- Claessen, D., Gilligan, C.A., Lutman, P.J.W., van den Bosch, F., 2005a. Which traits promote persistence of feral GM crops? Part 1: implications of environmental stochasticity. *Oikos*, 110: 20-29.
- Claessen, D., Gilligan, C.A., van den Bosch, F., 2005b. Which traits promote persistence of feral GM crops? Part 2: implications of metapopulation structure. *Oikos*, 110: 30-42.
- Codex Alimentarius, 2003. Codex principles and guidelines on foods derived from biotechnology. Joint FAO/WHO Food Standards Programme, Food and Agriculture Organisation, Rome.
- Colbach, H., 2009. How to model and simulate the effects of cropping systems on population dynamics and gene flow at the landscape level: example of oilseed rape volunteers and their role for co-existence of GM and non-GM crops. *Environmental Science and Pollution Research*, 16: 348-360.
- Colbach, N., Clermont-Dauphin, C., Meynard, J.M., 2001a. GeneSys: a model of the influence of cropping system on gene escape from herbicide tolerant rapeseed crops to rape volunteers I. Temporal evolution of a population of rapeseed volunteers in a field. *Agriculture, Ecosystems and Environment*, 83: 235-254.
- Colbach, N., Clermont-Dauphin, C., Meynard, J.M., 2001b. GeneSys: a model of the influence of cropping system on gene escape from herbicide tolerant rapeseed crops to rape volunteers II. Genetic exchanges among volunteer and cropped population a small region. *Agriculture, Ecosystems and Environment*, 83: 255-270.
- Colbach, N., Molinari, N., Meynard, J.M., Messéan, A., 2005. Spatial aspects of gene flow between rapeseed varieties and volunteers. *Agronomy for Sustainable Development*, 25: 355-368.
- Crawley, M.J., Brown, S.L., 1995. Seed limitation and the dynamics of feral oilseed rape on the M25 motorway. *Proceedings of the Royal Society B-Biological Sciences*, 259: 49-54.

- Crawley, M.J., Brown, S.L., 2004. Spatially structured population dynamics in feral oilseed rape. *Proceedings of the Royal Society B-Biological Sciences*, 271: 1909-1916.
- Crawley, M.J., Hails, R.S., Rees, M., Kohn, D., Buxton, J., 1993. Ecology of transgenic oilseed rape in natural habitats. *Nature*, 363: 620-623.
- Crawley, M.J., Brown, S.L., Hails, R.S., Kohn, D.D., Rees, M., 2001. Transgenic crops in natural habitats. *Nature*, 409: 682-683.
- Devaux, C., Lavigne, C., Falentin-Guyomarc'h, H., Vautrin, S., Lecomte, J., Klein, E.K., 2005. High diversity of oilseed rape pollen clouds over an agro-ecosystem indicated long-distance dispersal. *Molecular Ecology*, 14: 2269-2280.
- Devaux, C., Lavigne, C., Austerlitz, F., Klein, E.K., 2007. Modelling and estimating pollen movement in oilseed rape (*Brassica napus*) at the landscape scale using genetic markers. *Molecular Ecology*, 16: 487-499.
- Devaux, C., Klein, E.K., Lavigne, C., Sausse, C., Messéan, A., 2008. Environmental and landscape effects on cross-pollination rates observed at the long distance among French oilseed rape (*Brassica napus*) commercial fields. *Journal of Applied Ecology*, 45: 803-812.
- Devos, Y., Reheul, D., De Schrijver, A., Cors, F., Moens, W., 2004. Management of herbicide-tolerant oilseed rape in Europe: a case study on minimizing vertical gene flow. *Environmental Biosafety Research*, 3: 135-148.
- Devos, Y., De Schrijver, A., Reheul, D., 2009. Quantifying the introgressive hybridisation propensity between transgenic oilseed rape and its wild/weedy relatives. *Environmental Monitoring and Assessment*, 149: 303-322.
- Eastham, K., Sweet, J., 2002. Genetically modified organisms (GMOs): the significance of gene flow through pollen transfer, European Environment Agency, http://www.eea.europa.eu/publications/environmental_issue_report_2002_28
- EFSA, 2004. Opinion of the Scientific Panel on Genetically Modified Organisms on a request from the Commission related to the Notification (Reference C/NL/98/11) for the placing on the market of herbicide-tolerant oilseed rape GT73, for import and processing, under Part C of Directive 2001/18/EC from Monsanto. *The EFSA Journal*, 29: 1-19, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620772413.htm
- EFSA, 2005. Opinion of the Scientific Panel on Genetically Modified Organisms on a request from the Commission related to the application (Reference C/BE/96/01) for the placing on the market of glufosinate-tolerant hybrid oilseed rape Ms8 x Rf3, derived from genetically modified parental lines (Ms8, Rf3), for import and processing for feed and industrial uses, under Part C of Directive 2001/18/EC from Bayer CropScience. *The EFSA Journal*, 281: 1-23, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620770114.htm
- EFSA, 2006a. Guidance document of the Scientific Panel on Genetically Modified Organisms for the risk assessment of genetically modified plants and derived food and feed. *The EFSA Journal*, 99: 1-100, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620775747.htm

- EFSA, 2006b. Opinion of the Scientific Panel on Genetically Modified Organisms on the Post Market Environmental Monitoring (PMEM) of genetically modified plants. *The EFSA Journal*, 319: 1-27, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620769727.htm
- EFSA, 2008. Opinion of the Scientific Panel on Genetically Modified Organisms on an application (Reference EFSA-GMO-UK-2005-25) for the placing on the market of glufosinate-tolerant oilseed rape T45 for food and feed uses, import and processing and renewal of the authorization of oilseed rape T45 as existing products, both under Regulation (EC) 1829/2003 from Bayer CropScience. *The EFSA Journal*, 635: 1-22, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178690393760.htm
- Fredshavn, J.R., Poulsen, G., Huybrechts, I., Rüdelsheim, P., 1995. Competitiveness of transgenic oilseed rape. *Transgenic Research*, 4: 142-148.
- Garnier, A., Lecomte, J., 2006. Using spatial and stage-structured invasion model to assess the spread of feral population of transgenic oilseed rape. *Ecological Modelling*, 194: 141-149.
- Garnier, A., Deville, A., Lecomte, J., 2006. Stochastic modelling of feral plant populations with seed immigration and road verge management. *Ecological Modelling*, 197: 373-382.
- Garnier, A., Pivard, S., Lecomte, J., 2008. Measuring and modelling anthropogenic secondary seed dispersal along roadverges for feral oilseed rape. *Basic and Applied Ecology*, 9: 533-541.
- Gruber, S., Claupein, W., 2006. Fecundity of volunteer oilseed rape and estimation of potential gene dispersal by a practice-related model. *Agriculture, Ecosystems and Environment*, 119: 401-408.
- Gruber, S., Colbach, N., Barbottin, A., Pekrun, C., 2008. Post-harvest gene escape and approaches for minimizing it. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*, 3: 1-17.
- Hails, R.S., Bullock, J.M., Morley, K., Lamb, C., Bell, P., Horsnell, R., Hodgson, D.J., Thomas, J., 2006. Predicting fitness changes in transgenic plants: testing a novel approach with pathogen resistant Brassicas. *IOBC/WPRS Bulletin*, 29: 63-70.
- Jørgensen, R.B., 2007. Oilseed rape: Co-existence and gene flow from wild species. *Advances in Botanical Research*, 45: 451-464.
- Jørgensen, R.B., Hauser, T., D'Hertefeldt, T., Andersen, N.S., Hooftman, D., 2009. The variability of processes involved in transgene dispersal – case studies from *Brassica* and related genera. *Environmental Science and Pollution Research*, pre-print published online, DOI:10.1007/s11356-009-0142-4.
- Kawata, M., Murakami, K., Ishikawa, T., 2008. Dispersal and persistence of genetically modified oilseed rape around Japanese harbors. *Environmental Science and Pollution Research*, 16: 120-126.

- Knispel, A.L., McLachlan, S.M., Van Acker, R.C., Friesen, L.F., 2008. Gene flow and multiple herbicide resistance in escaped canola populations. *Weed Science*, 56: 72-80.
- Légère, A., 2005. Risks and consequences of gene flow from herbicide-resistant crops: canola (*Brassica napus* L) as a case study. *Pest Management Science*, 61: 292-300.
- Lecoq, E., Holt, K., Janssens, J., Legris, G., Pleysier, A., Tinland, B., Wandelt, C., 2007. General surveillance: roles and responsibilities the industry view. *Journal of Consumer Protection and Food Safety*, 2(S1): 25-28.
- Lutman, P.J.W., Berry, K., Payne, R.W., Simpson, E., Sweet, J.B., Champion, G.T., May, M.J., Wightman, P., Walker, K., Lainsbury, M., 2005. Persistence of seeds from crops of conventional and herbicide tolerant oilseed rape (*Brassica napus*). *Proceedings of the Royal Society B-Biological Sciences*, 272: 1909-1915.
- Lutman, P.J.W., Sweet, J., Berry, K., Law, J., Payne, R., Simpson, E., Walker, K., Wightman, P., 2008. Weed control in conventional and herbicide tolerant winter oilseed rape (*Brassica napus*) grown in rotations with winter cereals in the UK. *Weed Research*, 48: 408-419.
- Menzel, G., 2006. Verbreitungsdynamik und Auskreuzungspotential von *Brassica napus* L. (Raps) im Großraum Bremen. GCA-Verlag, Waabs, ISBN 3-89863-213-X.
- Messéan, A., Sausse, C., Gasquez, J., Darmency, H., 2007. Occurrence of genetically modified oilseed rape seeds in the harvest of subsequent conventional oilseed rape over time. *European Journal of Agronomy*, 27: 115-122.
- Neuffer, B., 2009. Kulturpflanzen als Quelle für Neophyten – Etablierungsmechanismen bei der Verwilderung von Raps (2004-2006), und Abschätzung des Hybridisierungspotentials ruderaler Raps- und Rübsenpopulationen (2007-2009) Im Rahmen des BIOLOG EUROPA Projektverbundes "Evolutionäre, ökologische und gesellschaftliche Konsequenzen biologischer Invasionen", http://www.gmo-safety.eu/en/safety_science/188.docu.html
- Nishizawa, T., Nakajima, N., Aono, M., Tamaoki, M., Kubo, A., Saji, H., 2009. Monitoring the occurrence of genetically modified oilseed rape growing along a Japanese roadside: 3-year observations. *Environmental Biosafety Research*, 8: 33-44.
- Norris, C., Sweet, J., 2002. Monitoring large scale releases of genetically modified crops (EPG1/5/84) incorporating report on project EPG 1/5/30: monitoring releases of genetically modified crop plants, DEFRA report, EPG 1/5/84, http://www.defra.gov.uk/environment/gm/research/pdf/epg_1-5-84_screen.pdf
- Pascher, K., Narendja, F., Rau, D., 2006. Feral oilseed rape – Investigations on its potential for hybridisation, Studie im Auftrag des Bundesministeriums fuer Gesundheit und Frauen, Forschungsberichte der Sektion IV, Band 3/2006, http://www.bmgfj.gv.at/cms/site/attachments/8/1/9/CH0255/CMS1138950978238/feral_oilseed_rape_-_investigation_on_its_potential_for_hybridisation_gesamt_f_hp.pdf

- Pessel, F.D., Lecomte, J., Emeriau, V., Krouti, M., Messéan, A., Gouyon, P.H., 2001. Persistence of oilseed rape (*Brassica napus* L.) outside of cultivated fields. *Theoretical and Applied Genetics*, 102: 841-846.
- Pivard, S., Adamczyk, K., Lecomte, J., Lavigne, C., Bouvier, A., Deville, A., Gouyon, P.H., Huet, S., 2008a. Where do the feral oilseed rape populations come from? A large-scale study of their possible origin in a farmland area. *Journal of Applied Ecology*, 45: 476-485.
- Pivard, S., Demšar, D., Lecomte, J., Debeljak, M., Džeroski, S., 2008b. Characterizing the presence of oilseed rape feral populations on field margins using machine learning. *Ecological Modelling*, 212: 147-154.
- Reuter, H., Menzel, G., Pehlke, H., Breckling, B., 2008. Hazard mitigation or mitigation hazard? Would genetically modified dwarfed oilseed rape (*Brassica napus*) increase feral survival? *Environmental Science and Pollution Research*, 15: 529-535.
- Saji, H., Nakajima, N., Aono, M., Tamaoki, M., Kubo, A., Wakiyama, S., Hatase, Y., Nagatsu, M., 2005. Monitoring the escape of transgenic oilseed rape around Japanese ports and roadsides. *Environmental Biosafety Research*, 4: 217-222.
- Scheffler, J.A., Dale, P.J., 1994. Opportunities for gene transfer from transgenic oilseed rape (*Brassica napus*) to related species. *Transgenic Research*, 3: 263-278.
- Simard, M.J., Légère, A., Pageau, D., Lajeunesse, J., Warwick, S., 2002. The frequency and persistence of canola (*Brassica napus*) volunteers in Québec cropping systems. *Weed Technology*, 16: 433-439.
- Simard, M.J., Légère, A., Séguin-Swartz, G., Nair, H., Warwick, S., 2005. Fitness of double vs. single herbicide-resistant canola. *Weed Science*, 53: 489-498.
- Spök, A., Hofer, H., Lehner, P., Valenta, R., Stirn, S., Gaugitsch, H., 2004. Risk assessment of GMO products in the European Union. Umweltbundesamt Wien. Berichte, Band 253. July 2004, http://www.umweltbundesamt.at/en/publikationen/publikationssuche/publikationsdetail/?pub_id=1531
- Spök, A., Gaugitsch, H., Laffer, S., Pauli, G., Saito, H., Sampson, H., Sibanda, E., Thomas, W., Van Hage, M., Valenta, R., 2005. Suggestions for the assessment of the allergenic potential of genetically modified organisms. *International Archives of Allergy and Immunology*, 137: 167-180.
- Sweet, J., Simpson, E., Law, J., Lutma, P., Berry, K., Payne, R., Champion, G., May, M., Walker, K., Wightman, P., Lainsbury, M., 2004. Botanical and Rotational Implications of Genetically Modified Herbicide Tolerance (BRIGHT) HGCA Project Report 353, 265. HGCA London, UK.
- von der Lippe, M., Kowarik, I., 2007a. Long-distance dispersal of plants by vehicles as a driver of plant invasions. *Conservation Biology*, 21: 986-996.
- von der Lippe, M., Kowarik, I., 2007b. Crop seed spillage along roads: a factor of uncertainty in the containment of GMO. *Ecography*, 30: 483-490.

- Warwick, S.I., Beckie, H.J., Small, E., 1999. Transgenic crops: new weed problems for Canada? *Phytoprotection*, 80: 71-84.
- Warwick, S., Beckie, H.J., Simard, M.J., Légère, A., Nair, H., Séguin-Swartz, G., 2004. Environmental and agronomic consequences of herbicide-resistant (HR) canola in Canada. In: den Nijs, H.C.M., Bartsch, D., Sweet, J. (Eds), *Introgression from Genetically Modified Plants into Wild Relatives* (pp 323-337), CABI publishing.
- Warwick, S.I., Simard, M.J., Légère, A., Beckie, H.J., Braun, L., Zhu, B., Mason, P., Séguin-Swartz, G., Stewart, C.N. Jr., 2003. Hybridization between transgenic *Brassica napus* L. and its wild relatives: *B. rapa* L., *Raphanus raphanistrum* L., *Sinapis arvensis* L., and *Erucastrum gallicum* (Willd.) O.E. Schulz. *Theoretical and Applied Genetics*, 107: 528-539.
- Warwick, S.I., Légère, A., Simard, M.-J., James, T., 2008. Do escaped transgenes persist in nature? The case of an herbicide resistance transgene in a weedy *Brassica rapa* population. *Molecular Ecology*, 17: 1387-1395.
- Wichmann, M.C., Alexander, M.J., Soons, M.B., Galsworthy, S., Dunne, L., Gould, R., Fairfax, C., Niggemann, M., Hails, R.S., Bullock, J.M., 2009. Human-mediated dispersal of seeds over long distances. *Proceedings of the Royal Society B-Biological Sciences*, 276: 523-532.
- Wilkinson, M.J., Timmons, A.M., Charters, Y., Dubbels, S., Robertson, A., Wilson, N., Scott, S., O'Brien, E., Lawson, H.M., 1995. Problems of risk assessment with genetically modified oilseed rape. In: *Proceedings of the Brighton Crop Protection Conference – Weeds*, 3: 1035-1044.
- Windels, P., Alcalde, E., Lecoq, E., Legris, G., Pleysier, A., Tinland, B., Wandelt, C., 2008. General surveillance for import and processing: the EuropaBio approach. *Journal of Consumer Protection and Food Safety*, 3(S2): 14-16.
- Yoshimura, Y., Beckie, H.J., Matsuo, K., 2006. Transgenic oilseed rape along transportation routes and port of Vancouver in western Canada. *Environmental Biosafety Research*, 5: 67-75.
- Zwaenepoel, A., Roovers, P., Hermy, M., 2006. Motor vehicles as vectors of plant species from road verges in a suburban environment. *Basic and Applied Ecology*, 7: 83-93.

SCIENTIFIC OPINION

Request from the European Commission related to the safeguard clause invoked by Austria on maize lines MON863 according to Article 23 of Directive 2001/18/EC¹

Scientific Opinion of the Panel on Genetically Modified Organisms

(Question No EFSA-Q-2008-742)

Adopted on 15 June 2009

PANEL MEMBERS*

Hans Christer Andersson, Salvatore Arpaia, Detlef Bartsch, Josep Casacuberta, Howard Davies, Patrick du Jardin, Lieve Herman, Niels Hendriksen, Sirpa Kärenlampi, Jozsef Kiss, Gijs Kleter, Ilona Kryspin-Sørensen, Harry Kuiper, Ingolf Nes, Nickolas Panopoulos, Joe Perry, Annette Pötting, Joachim Schiemann, Willem Seinen, Jeremy Sweet and Jean-Michel Wal.

SUMMARY

On 16 July 2008, Austria invoked Article 23 of Directive 2001/18/EC (safeguard clause) to provisionally prohibit the import of the MON863 maize lines on its territory. Austria provided detailed reasons listed in supporting documents.

On 3 September 2008, the European Food Safety Authority (EFSA) has been requested by the European Commission to provide a scientific opinion on the statement and documents submitted by Austria in the context of the safeguard clause invoked under Article 23 of Directive 2001/18/EC.

In the light of the information package provided by Austria in support of its safeguard clause and, having considered all relevant scientific publications, the Scientific Panel on Genetically Modified Organisms (GMO Panel) of EFSA concludes that, in terms of risk to human and animal health and the environment, no new scientific evidence was presented that would invalidate the previous risk assessment of maize MON863. The EFSA GMO Panel also concludes that no new scientific data or information was provided in support of adverse effects of maize MON863 on the environment and on human and animal health in Austria.

¹ For citation purposes: Scientific Opinion of the Panel on Genetically Modified Organisms on a request from the European Commission related to the safeguard clause invoked by Austria on maize lines MON863 according to Article 23 of Directive 2001/18/EC. *The EFSA Journal* (2009) 1152, 1-18.

* (minority opinion) This opinion is not shared by x members of the Panel. / (conflict of interest) x members of the Panel did not participate in (part of) the discussion on the subject referred to above because of possible conflicts of interest.

Therefore, no specific scientific evidence, in terms of risk to human and animal health and the environment, were provided that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC.

Key words: GMOs, maize (*Zea mays* L.), MON863, Austria, safeguard clause, human health and animal health, environment, Directive 2001/18/EC

TABLE OF CONTENTS

Panel Members.....	1
Summary	1
Table of Contents	3
Background	4
Terms of reference as provided by the European Commission	4
Acknowledgements	4
Assessment	5
1. Introduction	5
2. Assessment of documents provided by Austria	5
2.1. Food and feed safety issues.....	6
2.1.1. Toxicological risk assessment.....	6
2.1.2. Nutritional risk assessment	
2.2. Molecular characterisation issues	8
2.3. Environmental safety issues and post-market environmental monitoring8	
2.3.1. Environmental risk assessment	11
2.3.2. Post-market environmental monitoring plan.....	12
2.3.3. Conclusion	13
Overall Conclusions and Recommendations.....	13
Documentation provided to EFSA	14
References	14

BACKGROUND

On 16 July 2008, Austria notified to the European Commission a national safeguard clause on genetically modified (GM) maize MON863 under Article 23 of Directive 2001/18/EC. The notification was accompanied by the scientific document entitled “*Scientific arguments for an import ban of genetically modified maize MON863 (Zea mays L., MON863) of Monsanto notification C/DE/02/9*”.

On 3 September 2008, the European Food Safety Authority (EFSA) has received a request from the European Commission to provide a scientific opinion from its Scientific Panel on Genetically Modified Organisms (EFSA GMO Panel) on the statement and documents submitted by Austria in the context of its invoked safeguard clause.

TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

EFSA was requested, under Article 29(1) and in accordance with Articles 22(2) and 22(5)(c) of Regulation (EC) No 178/2002, to provide a scientific opinion as to “*whether, in accordance with Article 23 of Directive 2001/18/EC, the statement and documents submitted by the Austrian authorities comprise new information affecting the environmental risk assessment, such that detailed grounds exist to consider that the above authorised GMO, for the uses laid down in the corresponding consent, constitute a risk to human health, animal health or the environment*”.

ACKNOWLEDGEMENTS

The European Food Safety Authority wishes to thank the members of the Working Groups on Molecular Characterisation, Food/Feed and Environment, as well as the following members of its staff: Jaime Aguilera, Yann Devos, Yi Liu and Sylvie Mestdagh for the preparation of this opinion.

ASSESSMENT

1. Introduction

Directive 2001/18/EC provides the possibility for Member States to invoke safeguards on specific Genetically Modified Organisms (OGM) in the case where new or additional information, made available since the date of the consent, would affect the risk assessment of an authorised GMO. Provisions foreseen by Austria seek to provisionally prohibit the import of maize lines MON863 into the Austrian territory.

The EFSA GMO Panel examined the set of supporting documents submitted by Austria. In this respect, the EFSA GMO Panel assessed whether the submitted documents comprise new scientific knowledge information that would change the outcome of previously performed risk assessment, and if detailed grounds exist to consider that the authorised maize MON863, for its intended uses, constitutes a risk to human and animal health or the environment.

The EFSA GMO Panel looked for evidence for GMO-specific risks – including long-term effects (e.g., BEETLE report, 2009) – taking into consideration the EFSA GMO Panel guidance for the risk assessment of genetically modified plants and derived food and feed (EFSA, 2006a) as well as any related risk assessments carried out in the past. In addition, the EFSA GMO Panel considered the relevance of raised concerns in the light of the most recent scientific data and relevant peer-reviewed publications.

2. Assessment of documents provided by Austria

A set of supporting documents, accompanying the mandate of the European Commission (see *Terms of Reference as provided by the European Commission*), was forwarded to EFSA on 3 September 2008.

Austria provided the following set of documents in support of its safeguard clause:

- Verbot des Inverkehrbringens von gentechnisch verändertem Mais der Linie MON863;
- Scientific arguments for an import ban of maize MON863.

Based on the supporting documents, several issues were identified and therefore considered by the EFSA GMO Panel in the following areas: (1) the toxicological and nutritional risk assessment, (2) the safety assessment of the antibiotic resistance marker (ARM) gene *npII*, and (3) the environmental risk assessment and post-market environmental monitoring plan related to the accidental spillage of maize MON863.

In addition, the EFSA GMO Panel notes that it only gives its opinion on the scientific quality of the post-market environmental monitoring plan proposed by the applicant, whilst the final endorsement thereof is done by risk managers.

During its assessment, the EFSA GMO Panel identified issues raised by Austria that would require further clarifications. To present and clarify the provided set of documents, an informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel

and EFSA staff was held on 23 April 2009. A representative of the European Commission attended the meeting as observer.

2.1. Food and feed safety issues

The Austrian Competent Authority has provided EFSA with a document entitled “Scientific Arguments for an Import Ban of Genetically Modified Maize MON 863 (*Zea mays* L., line MON 863) of Monsanto (Notification C/DE/02/9)”. This document focuses on a number of details of toxicological and nutritional studies with maize MON863, which are cited by an article by Hammond et al. (2006) describing the outcomes of the 90-day rat oral toxicity study with maize MON863. The study by Hammond et al. (2006) refers to the same study for which another technical report was previously provided to EFSA as part of the dossier that was considered by the EFSA GMO Panel when formulating its opinion. In addition, various other documents have been published by EFSA in response to issues previously raised on the same 90-days study being part of the dossier (EFSA, 2004a; EFSA, 2004b; EFSA, 2007a; EFSA, 2007b). None of the studies quoted by Austria provide new direct evidence of risks to human and animal health associated with food and feed use of MON863, or its transgenic components, which would have evolved after the publication of EFSA’s opinion on maize MON863 in 2004. In addition, the EFSA GMO Panel notes that 90-day study is not strictly required in the internationally harmonized approach of comparative safety assessment of a GMO versus its control. If the outcomes of the comparison of extensively analyzed compositional and other parameters indicate equivalence of a GMO to conventional food and feed, this would not trigger the performance of a 90-days study.

2.1.1. Toxicological risk assessment

The Austrian Competent Authority highlighted a number of details of the toxicological and nutritional studies cited by Hammond et al. (2006) as what they consider as deficiencies in experimental design and interpretation, which, as they assert, would impact on the interpretability of the outcomes of these studies. With regard to toxicity, the Austrian comments focus on the 90-days feeding study with diets containing kernels of MON863. They highlight various perceived deficiencies in the original dossier report on the 90-days study.

As the Austrian Competent Authority correctly notes, a report on the same 90-days oral rat feeding study as described by Hammond et al. (2006) has been provided with the dossier on MON863, albeit with more extensive coverage of experimental details.

With regard to the 90-days rat feeding study, the Austrian Competent Authority has made detailed and extensive comments regarding the experimental design (no updated protocol, inclusion of reference groups), diet formulation (Cry3Bb denaturation; contaminant analysis), endpoints measured (not all organs/tissues checked for histopathology; various single differences in performance and clinical chemistry). The same data of this study have already been assessed in detail by the EFSA GMO Panel, including the various differences observed (EFSA, 2004a; EFSA, 2004b; EFSA, 2007a; EFSA, 2007b)

Various general comments can be made, though, to the conclusions and assertions brought forward in the Austrian document. The Austrian Competent Authority, for example, appears to comment on the statistical analysis as performed by the applicant, including the asserted disappearance of differences through an increase of the total number of animals used for the

experiment, most of which comprise animals in reference groups fed diets containing commercial maize. The EFSA GMO Panel, though, has considered the direct comparison between the groups fed GM maize MON863 and the respective control groups. Any difference thus identified can then be further compared with the background ranges provided by the animals fed the reference diets. For example, the difference in reticulocytes that the Austrian authorities point at, was also noted by the EFSA GMO Panel, as summarized in section 4.2 of its opinion of 2004 (EFSA, 2004a). This approach of direct comparison of a GMO with its appropriate comparator is in line with the internationally harmonized comparative safety assessment principles for GM foods as laid down in Codex alimentarius guidelines (to which the Austrian authorities have also subscribed) and further elaborated in EFSA guidance document (EFSA, 2006a). Concerning the comment with regard to the percentage (60-80%) of the reference group in overall sample size, it should be noted that it is not the percentage of the reference groups within the sample size that is a problem in itself, but the replication of the GM and comparator. If this replication is too small because of the restrictions placed by having 60-80% reference varieties, then this could be a valid criticism. However, if the replication is sufficient, then there is no reason why reference groups should not contribute 60-80% of the sample size. In addition, the guidelines that were recently drafted by the Commission and discussed with EU Member States' representatives at the Standing Committee for the Food Chain and Animal Health recommend that compositional field trials be done with at least 60% of the plots being allocated to commercial varieties (reference groups).

Austria also refers to OECD guideline 408 (OECD, 1998) with regard to the number of doses and other details, which would not have been strictly followed by the applicant. In the EFSA GMO Panel's opinion, it has to be taken into account, though, that OECD guideline 408 has been developed for the testing of pure chemicals and that adjustments may be required for the testing of whole foods. This issue has also been extensively highlighted in the EFSA GMO Panel's report on animal feeding trials (EFSA, 2008).

In addition, Austria notes that the Cry3Bb protein should have remained intact in the test diet used in the 90-days rat feeding study, and therefore concludes that more details on the feed manufacturing conditions and the presence and nature of the transgenic elements in the rat diets should have been provided. Given the commonly low expression levels of transgenic proteins in GM crops, the EFSA GMO Panel notes, though, that for testing the safety of transgenic proteins, other models than whole-product feeding studies for testing purified components would be recommendable, such as repeated-dose oral rodent studies with the purified transgenic protein (see EFSA guidance document, EFSA, 2006a). Based on the data provided in the dossier, Cry3Bb has been assessed by the EFSA GMO Panel for its safety. The EFSA GMO Panel concludes that the data provided by the Austrian Competent Authority do not indicate that there would be an additional need for further testing the safety of Cry3Bb.

The EFSA GMO Panel concurs with the Austrian notion that nutritional feeding studies in food-producing animals cannot be considered toxicity studies. Again, the Austrian comments pertain to conclusions made by other authors than the EFSA GMO Panel in its opinion on MON863. The scientific opinions of the EFSA GMO Panel, including the one on MON863, make a distinction between toxicity, allergenicity, nutrition, and other issues that may be of relevance to food and feed safety.

In conclusion, no new data have been presented by the Austrian authorities that could be considered evidence of potential toxic effects on maize MON863 and its transgenic components on humans and animals. In the absence of such evidence, the EFSA GMO Panel

cannot follow the Austrian recommendation for requiring additional toxicity tests for sub-chronic, chronic, developmental, and reproductive toxicity of maize MON863.

2.1.2. Nutritional risk assessment

The Austrian Competent Authority refers to various nutritional studies that have appeared in scientific journals, in which the performance of various food-producing animals fed MON863, including poultry, beef and dairy cattle, and swine, have been measured. The article on the poultry feeding study with MON863 (Taylor et al., 2003) is based on the same feeding trial of which a report has been provided with the dossier assessed by EFSA. In addition, the quoted article on the dairy cattle study (Grant et al., 2003) has also been considered in the EFSA opinion. The other articles describing studies on beef cattle and swine were published after the publication of the opinion.

The Austrian comments pertain to perceived shortcomings of the abovementioned nutritional studies, such as to the composition of the diets (e.g., origin of the maize; certain micronutrients not being measured; levels of transgenic components in final product), experimental design (e.g., inclusion of reference groups; choice of animals; background variability), and statistical analysis (e.g. limited information based on p-values alone). None of these comments indicate that the new data cited by Austria demonstrate a potentially relevant adverse effect of maize MON863 on animal nutrition.

The EFSA GMO Panel concurs with the Austrian general comment that the first comparison to be made is between the GMO and its appropriate comparator, after which further comparison with the background range of reference values may be made, if applicable. The Austrian Competent Authority assert that whilst the transgenic identity of test materials has been verified, the levels of transgenic proteins and DNA in the final feed product, rather than in raw products, should be established. In the EFSA GMO Panel's opinion, these transgenic components do not contribute to nutrition given their very low abundance.

Moreover, the Austrian comments and the cited studies do not provide evidence of potential impacts of maize MON863 on nutrition that would lead the EFSA GMO Panel to deviate from its previous opinion (EFSA, 2004a). It should be noted that this kind of studies is not strictly required in the absence of indications of altered nutritional properties of a GMO based on compositional analysis and other data (e.g. type of modification) according to EFSA guidance. In the case of maize MON863, these data can therefore be regarded as supplementary data.

2.2. Molecular characterisation issues

2.2.1. Safety assessment of the ARM gene *aph(3')*-IIa

Therapeutic relevance of kanamycin and neomycin in human and veterinary medicine

The *aph(3')*-IIa gene confers resistance to neomycin and kanamycin but not to other aminoglycosides of clinical use. The *aph(3')*-IIa gene confers slightly reduced susceptibility to amikacin for *E. coli*. However, amikacin is a poor substrate for the APH(3')-IIa enzyme due to its hydroxyaminobutyryl side chain.

Antimicrobials are grouped into classes on the basis of chemical structure and mode of action. Most antimicrobials used for the treatment of animals belong to classes that are also used in

human medicine. A list of antimicrobial classes was compiled by WHO in 2007 (the WHO Expert Group on Critically Important Antimicrobials for Human Health) (WHO, 2007), giving examples of substances used for the treatment of infections in humans. Antimicrobials listed as ‘Critically Important’ in the WHO list are characterised as both: 1. ‘sole therapy or one of few alternatives to treat serious human disease’, and 2. ‘antimicrobials used to treat diseases caused by organisms that may be transmitted *via* non-human sources or diseases caused by organisms that may acquire resistance genes from non-human sources’. With respect to this classification, kanamycin and neomycin were downscaled from “critically important” to “highly important” compared to the WHO report of 2005 (WHO, 2005), because they no longer were considered to comply with criterion 1. Antimicrobials classified as ‘Highly Important’ meet either criterion 1 or criterion 2. The World Animal Health Organisation (OIE) has similarly developed and adopted a list ranking the importance of different antimicrobials for animal health (OIE, 2007). Aminoglycosides, as a group, is a class of antibiotics critically important for veterinary medicine and animal production (OIE, 2007).

Kanamycin is rarely used systemically today due to its severe side effects although this antibiotic remains a recognised second line choice in conditions of infections with multiple drug resistant (MDR) *Mycobacterium tuberculosis* (MTB). Aminoglycoside resistance in *M. tuberculosis* results from a mutation causing alterations in the antibiotic target molecule within the mycobacterial cell; thus chromosomal resistance, and not the transfer of antibiotic resistance genes [such as *aph(3’)-IIa*], is the only identified mechanism resulting in resistance to kanamycin (Goldstein *et al.*, 2005 and references therein). Neomycin is poorly absorbed from the gastrointestinal tract, and is nephrotoxic and ototoxic. The use of neomycin in human medicine is limited to topical applications and gut irrigation/encephalopathy. By killing bacteria in the intestinal tract, it keeps ammonia levels low and prevents hepatic encephalopathy, especially prior to gastrointestinal surgery. In veterinary medicine kanamycin and neomycin could be used for therapies of neonatal diarrhoea in piglets and treatment of multi-resistant enteric gram-negative infections.

Transfer of ARM genes from GM plants to bacteria

Transfer of antibiotic resistance marker genes from GM plants to bacteria has not been detected either under natural or laboratory conditions in the absence of pre-existing sequence identity in the recipient organism. A number of studies have been published in which the possible occurrence of bacterial transformants carrying GM plant-derived antibiotic resistance marker genes in fields planted with GM plants were screened (Paget *et al.*, 1998; Gebhard and Smalla, 1999; Badosa *et al.*, 2004; Demanèche *et al.*, 2008, Kim *et al.*, 2008). Nonetheless, in none of these studies has transfer of antibiotic resistance marker genes from GM plant material to bacteria been demonstrated. Nor could such transfer be detected from the existing background.

In the cascade of events leading to clinical importance, the ARM genes present in GM plants would need to be transferred to, stabilised and expressed in a bacterial cell. Absence of sequence identity is known to be the major factor limiting the stable integration, by homologous recombination, of DNA from a GM plant to a bacterial cell. Other factors limiting the potential impact of the *aph(3’)-IIa* marker gene used in maize MON863 include the lack of stability of the plant DNA in different environments and the limited competence of many bacterial cells.

ARM genes transfer from plants to bacteria has never been observed. There are limitations related to sampling and detection methods, as well as challenges in estimating exposure levels and in the inability to assign transferable resistance genes to a defined source. Notwithstanding these uncertainties, and taking into account all the peer-reviewed data, the probability of the transfer of antibiotic resistance genes from GM plants to bacteria must be considered as zero or below the limit of detection. It has to be stressed that the dissemination of antibiotic resistance genes between bacteria is not comparable with the potential transfer of these genes from plants to bacteria. Transfer of ARM genes from plants to bacteria would occur, if at all, with a very low frequency compared to that between bacteria.

Resistance in natural environments

Concerning the prevalence of the *aph(3')*-IIa gene and phenotypic kanamycin/neomycin resistance, two basically different sets of data are available.

In different countries the resistance is monitored in indicator bacteria and pathogens. The available data indicate that a pool of *aph(3')*-IIa-carrying bacteria occur in and outside hospital-associated environments. There are environmental fluctuations in the prevalence of the *aph(3')*-IIa gene and of the kanamycin/neomycin phenotypic resistance. During the bilateral technical meeting between the GMO Panel and the Austrian delegation a mean frequency of 0.74 % kanamycin resistant *Salmonella* has been reported in the Austrian surveillance programme of human isolates from 1999-2008. A similar frequency has been reported for neomycin resistant *Campylobacter* isolates from poultry from 2004-2007. This should be considered as high frequency.

An additional and important source of information on the frequency and distribution of genes conferring resistance to aminoglycoside antibiotics has been generated from metagenomic analyses. These are based on the molecular detection of resistance determinants, and allow the analysis of an extended bacterial population compared to conventional culturing methods. The expanding metagenomic studies have revealed a high density of antibiotic resistance genes in the environment (D'Costa et al., 2007). The resistance mechanisms identified include inactivation of aminoglycoside antibiotics by phosphorylation and acetylation (Riesenfeld et al., 2004). The results indicate that soil bacteria are a reservoir of antibiotic resistance genes towards aminoglycosides with greater genetic diversity than previously accounted for. Even a remote Alaskan soil, with no known exposure to antibiotics, harbours a great variety of resistance determinants (Allen et al., 2009) and even before the clinical use of antibiotics, antibiotic resistant bacteria were isolated (Wright, 2007). The ubiquitous distribution of the wide variety of antibiotic resistance genes can be explained by the fact that many of these genes are not just weapons against bacterial competitors but have other primary signalling functions (e.g. detoxification of metabolic intermediates, virulence and signal trafficking) (Martinez, 2008).

Selective antibiotic pressure in field environments

The EFSA GMO Panel agrees that selective antibiotic pressure can be present in several environments. This selective pressure is the key factor in the selection of resistance from the environmental reservoir and in the dissemination of antibiotic resistance between bacteria. However, this does not increase the likelihood of transfer of ARM genes from GM plants to bacteria. In the unlikely event that such a transfer to a bacterial cell occurs, it would not add

to the existing pool of naturally occurring resistance genotypes present in the microbial resistome.

2.2.2. Conclusion

Kanamycin and neomycin are categorised by the WHO Expert Group on Critically Important Antimicrobials for Human Health as ‘Highly Important’ antimicrobials. Kanamycin is used as a second-line drug for the treatment of infections with multiple drug-resistant tuberculosis (MTB). The increasing occurrence worldwide of “extensively drug-resistant” isolates of MTB with resistance to second-line antibiotics such as kanamycin is a cause for global concern. The *aph(3’)-IIa* gene has not been implicated in such resistance.

The transfer of ARM genes from GM plants to bacteria has never been shown to occur under laboratory or natural conditions in the absence of sequence identity. If transfer of ARM genes from GM plants to bacteria occurs at all, its frequency is below the limit of detection. The process is therefore considered unlikely to impact on the occurrence of antibiotic resistance in humans, animals and the environment.

The genetic determinants conferring resistance to kanamycin and neomycin are detected in the environmental resistome and in all environments analysed by metagenomic analyses. They are also widespread among bacteria in different environments. Data provided from Austria indicate the spread of this resistance in human food-borne pathogens at a frequency of about 1%. Taking into account the theoretical transfer frequencies of ARM genes from GM plants to bacteria, in the unlikely event of the transfer of the *aph(3’)-IIa* gene from maize MON863 to a bacterium, its contribution to the existing pool of kanamycin resistance in bacteria would be negligible.

For further details concerning the safety assessment of the ARM gene *aph(3’)-IIa*, the EFSA GMO Panel refers to the EFSA statement (EFSA, 2009).

2.3. Environmental safety issues and post-market environmental monitoring

2.3.1. Environmental risk assessment

The intended uses of maize MON863 specifically exclude cultivation, so the environmental exposure is mainly limited to exposure through manure and faeces from the gastrointestinal tracts mainly of animals fed maize MON863, as well as to accidental release into the environment of MON863 grains during transportation and processing and subsequently to potential occurrence of sporadic feral plants.

Maize is highly domesticated and generally unable to survive in the environment without cultivation. Maize plants are not winter hardy in most regions of Europe: they have lost their ability to release seeds from the cob and do not occur outside cultivated land or disturbed habitats in agricultural landscapes of Europe, despite cultivation for many years. In addition, there are no cross-compatible wild relatives in Europe, and gene flow via pollen is largely restricted to neighbouring maize crops.

Maize MON863 has been developed to provide protection against certain coleopteran pests such as corn rootworms (*Diabrotica* spp.) by the introduction of a variant *Bacillus thuringiensis cry3Bb1* gene. The insect resistance trait can only be regarded as providing a selective advantage for the GM maize in cultivation under corn rootworm infestation

conditions. However, survival of maize outside of cultivation in Europe is mainly limited by a combination of low competitiveness, absence of a dormancy phase, susceptibility to diseases and to cold climate conditions. Since these general characteristics of maize MON863 are unchanged, the inserted insect resistance trait is not likely to provide a selective advantage outside of cultivation in Europe. Therefore, it is considered very unlikely that volunteers of this GM maize, or its progeny, will differ from conventional maize varieties in their ability to survive until subsequent seasons, or to establish feral populations under European environmental conditions. Since studies in Europe and elsewhere with maize MON863 have shown no altered survival, multiplication or dissemination characteristics except in the presence of the specific target organisms, the EFSA GMO Panel is of the opinion that the likelihood of unintended environmental effects as a consequence of spread of genes from this maize due to accidental spillage will not differ from that of conventional maize varieties.

Considering the intended uses of maize MON863, the environmental risk assessment is concerned with indirect exposure through manure and faeces from the gastrointestinal tracts mainly of animals fed maize MON863. In its previous environmental risk assessment of maize MON863 (EFSA, 2004a), the EFSA GMO Panel also considered the possibility that gene products, particularly Cry proteins might enter the environment either from the intestinal tracts of animals or through horizontal gene flow to bacteria. Data supplied by the applicant and other data published in the scientific literature suggested that most proteins would be degraded by enzymatic activity in the intestinal tract (Lutz et al., 2005; Wiedemann et al., 2006), so that a small amount of Cry protein would remain intact to pass out in faeces. Subsequently, there would be further degradation of proteins in the manure due to microbial processes. Even though it has been observed that Cry proteins can bind to certain soil particles (e.g., humic acids, clays) resulting in protection from degradation, a number of studies revealed that there is no accumulation of Cry proteins from GM crops in soil (Herman et al., 2001, 2002; Head et al., 2002; Baumgarte and Tebbe, 2005; Hopkins and Gregorich, 2005; Ahmad et al., 2005; Dubelman et al., 2005; Icoz and Stotzky, 2007; Krogh and Griffiths, 2007; Lawhorn et al., 2009).

Having considered the different routes of exposure to the environment, the EFSA GMO Panel reiterates its previous opinion that the likelihood of unintended environmental effects as a consequence of spread of genes from this maize will not differ from that of conventional maize varieties (EFSA, 2004a). The EFSA GMO Panel concludes that the Austrian submission provided no new scientific data or information in support of an adverse effect of maize MON863 on the environment and that would justify a national safeguard measure concerning this product.

2.3.2. Post-market environmental monitoring plan

Austria questioned the adequacy, relevance and completeness of the post-market environmental monitoring (PMEM) plan provided by the applicant for maize MON863. The EFSA GMO Panel notes that it gives its opinion on the scientific quality of the PMEM plans proposed by applicants. The definitive and final endorsement of post-market environmental monitoring is done by risk managers. In this context, the EFSA GMO Panel refers to the section 5.2 of its scientific opinion on post-market environmental monitoring of GM plants (EFSA, 2006b) stating that *'Details of the specific plans and methods of monitoring in each country should not be included in the original application. The GMO Panel advises that the application should describe the general approaches and methods that the applicant would apply in different commercialisation sites, including the type of dialogue that would be*

established with risk managers in each Member State. The implementation of general surveillance data collection at a regional and national scale will be dependent on the local circumstances prevailing at the time the consent is given. Thus detailed local arrangements will be developed by the applicant after the application has been accepted and will depend on where the crop will be grown, the scale of commercialisation, the nature of the cultivation systems and a range of other factors. Applicants are encouraged to establish contacts with national competent authorities at an early stage in the commercialisation planning.'

In its initial scientific opinion for the placing on the market of insect protected GM maize MON863 and MON863xMON810 (EFSA, 2004a), the EFSA GMO Panel concluded that the scope of the monitoring plan provided by the applicant was in line with the intended uses for the GMO since the environmental risk assessment did not cover cultivation. In addition, the EFSA GMO Panel advised that appropriate management systems should be in place to prevent seeds of the GM maize entering cultivation, as the latter requires specific approval under Directive 2001/18/EC.

The association of European traders, importing cereals, oilseeds and feedstuffs (Coceral) as well as the association of European port silos (Unistock) have recently joined with the European Association of Bioindustries (EuropaBio) in developing monitoring systems for GM maize grains imported through the main ports of entry and processing facilities in Europe (e.g., Spain, Portugal, Italy, the Netherlands). These monitoring systems are based on Hazard Analysis and Critical Control Points (HACCP) principles. Therefore, it is anticipated that risk managers will opt for post-market environmental monitoring of imported maize MON863 in accordance with these arrangements.

Subject to post-market environmental monitoring activities under the coordinating system established by EuropaBio, the EFSA GMO Panel maintains its opinion that the scope of the post-market environmental monitoring plan provided by the applicant complies with (1) the intended uses of maize MON863, which exclude cultivation, (2) the requirements of the EFSA GMO Panel guidance on GM plants (EFSA, 2006a), and (3) the EFSA GMO Panel scientific opinion on post-market environmental monitoring (EFSA, 2006a,b).

2.3.3. Conclusion

The EFSA GMO Panel confirms its opinion that the likelihood of unintended environmental effects as a consequence of spread of (trans)genes from maize MON863 will not differ from that of conventional maize varieties in the context of its intended uses. The EFSA GMO Panel agrees with the monitoring plan submitted by the applicant, especially now that comprehensive arrangements have been made by applicants and operators for monitoring at major points of import and processing in European Union. However, the EFSA GMO Panel continues to recommend that appropriate management systems should be in place to minimise accidental loss and spillage of transgenic maize grains during transportation, storage and handling in the environment, and processing into derived products.

OVERALL CONCLUSIONS AND RECOMMENDATIONS

The EFSA GMO Panel has investigated the claims and documents both submitted in support of the Austrian safeguard clause and presented at the informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel and EFSA staff on 23 April 2009. In

these documents, the EFSA GMO Panel did not identify any new data subject to scientific scrutiny or scientific information that would invalidate the previous risk assessment of maize MON863. In addition, the Austrian submission did not supply scientific evidence, that the environment or ecology of Austria was different from other regions of the EU, sufficient to merit separate risk assessments from those conducted for other regions in the EU.

Having considered the overall information package submitted by Austria as well as relevant scientific literature, the EFSA GMO Panel is of the opinion that there is no specific scientific evidence, in terms of risk to human and animal health and the environment, that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC for the import of maize MON863 for its intended uses in Austria. In conclusion, the EFSA GMO Panel finds that the scientific evidence currently available does not sustain the arguments provided by Austria, and therefore the EFSA GMO Panel reiterates its previous scientific opinions on maize MON863.

DOCUMENTATION PROVIDED TO EFSA

1. Letter, dated 3 September 2008, with supporting documents from Jos Delbeke, Acting Director-General Environment EC, to Catherine Geslain-Lanéelle, Executive Director EFSA (ref ENV/B3/YK/lh ARES(2008) 24354) requesting for a scientific opinion on the safeguard notification submitted by Austria under Article 23 of Directive 2001/18/EC for maize lines MON863 and comprising the following supporting documents:
 - a. Verbot des Inverkehrbringens von gentechnisch verändertem Mais der Linie MON863;
 - b. Scientific arguments for an import ban of maize MON863.
2. Letter, dated 15 October 2008, from Catherine Geslain-Lanéelle, Executive Director EFSA, to Jos Delbeke, Acting Director-General Environment EC, acknowledging the receipt of the mandate accompanied with the supporting documents.

REFERENCES

- Ahmad, A., Wilde G.,E., Zhu K., Y., 2005. Detectability of Coleopteran-specific Cry 3Bb1 protein in soil and its effect on nontarget surface and below-ground arthropods. *Environmental Entomology*, 34, 385-394.
- Allen, J.K., Moe, L.A., Rodbummer, J., Gaarder, A. and Handelsman, J., 2009. Functional metagenomics reveals diverse β -lactamases in a remote Alaskan soil. *International Society for Microbial Ecology*, 3, 243-251.
- Badosa, E., Moreno, C. and Montesinos, E., 2004. Lack of detection of ampicillin resistance gene transfer from Bt176 transgenic corn to culturable bacteria under field conditions. *FEMS Microbiol Ecology*, 48, 169-178.
- Baumgarte, S., Tebbe, C.C., 2005. Field studies on the environmental fate of the Cry1AB Bt toxin produced by transgenic maize (MON810) and its effect on bacterial communities in the maize rhizosphere. *Molecular Ecology*, 14, 2539-2551.
- BEETLE report, 2009. Long-term effects of genetically modified (GM) crops on health and the environment (including biodiversity): prioritization of potential risks and delimitation

- of uncertainties. German Federal Office of Consumer Protection and Food Safety, BLaU-Umweltstudien and Genius GmbH, http://ec.europa.eu/environment/biotechnology/pdf/beetle_report.pdf
- D'Costa, V.M., Griffiths, E. and Wright, G.D., 2007. Expanding the soil antibiotic resistome: exploring environmental diversity. *Current Opinion in Microbiology*, 10, 481-489.
- Demanèche, S., Sanguin, H., Poté, J., Navarro, E., Bernillon, D., Mavingui, P., Wildi, W., Vogel, T.M. and Simonet, P., 2008. Antibiotic-resistant soil bacteria in transgenic plant fields. *Proceedings of the National Academy of Sciences USA*, 105, 3957-3962.
- Dubelman, S., Ayden, B., Bader, B., Brown, C., Jiang, C., Vlachos, D., 2005. Cry1Ab protein does not persist in soil after 3 years of sustained Bt corn use. *Environmental Entomology*, 34, 915-921.
- EC, 2001. Directive 2001/18/EC of the European Parliament and of the Council of 12 March 2001 on the deliberate release into the environment of genetically modified organisms and repealing Council Directive 90/220/EEC. Official Journal of the European Communities, L106, 1-39, http://europa.eu.int/eur-lex/pri/en/oj/dat/2001/l_106/l_10620010417en00010038.pdf
- EFSA, 2004a. Opinion of the Scientific Panel on Genetically Modified Organisms on a request from the Commission related to the Notification (Reference C/DE/02/9) for the placing on the market of insect protected genetically modified maize MON863 and MON863xMON810, for import and processing, under Part C of Directive 2001/18/EC from Monsanto. The EFSA Journal 49, 1-25. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620772339.htm
- EFSA, 2004b. Statement on an Evaluation of the 13-week Rat Feeding Study on MON 863 Maize, submitted by the German Authorities to the European Commission by the Scientific Panel on Genetically Modified Organisms (GMO). European Food Safety Authority, Parma. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620775579.htm
- EFSA, 2006a. Guidance document of the Scientific Panel on Genetically Modified Organisms for the risk assessment of genetically modified plants and derived food and feed. The EFSA Journal, 99, 1-100, [including draft document updated in 2008. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1211902010430.htm], http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620775747.htm
- EFSA, 2006b. Opinion of the Scientific Panel on Genetically Modified Organisms on the Post Market Environmental Monitoring (PMEM) of genetically modified plants. The EFSA Journal, 319, 1-27. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620769727.htm
- EFSA, 2007a. Statement on the Analysis of Data from a 90-Day Rat Feeding Study with MON 863 Maize by the Scientific Panel on Genetically Modified Organisms (GMO). European Food Safety Authority, Parma. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178621169104.htm

- EFSA, 2007b. EFSA Review of Statistical Analyses conducted for the Assessment of the MON 863 90-Day Rat Feeding Study. European Food Safety Authority, Parma.
http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178621342614.htm
- EFSA, 2008. Safety and nutritional assessment of GM plants and derived food and feed: The role of animal feeding trials. Reports of the EFSA GMO Panel working group on animal feeding trials. *Food and Chemical Toxicology*, 46, S2-S70.
- EFSA, 2009. Statement of EFSA on the consolidated presentation of the joint Scientific Opinion of the GMO and BIOHAZ Panels on the “Use of Antibiotic Resistance Genes as Marker Genes in Genetically Modified Plants” and the Scientific Opinion of the GMO Panel on “Consequences of the Opinion on the Use of Antibiotic Resistance Genes as Marker Genes in Genetically Modified Plants on Previous EFSA Assessments of Individual GM Plants”. *The EFSA Journal*, 1108: 1-8,
http://www.efsa.europa.eu/EFSA/ScientificPanels/GMO/efsa_locale-1178620753812_Statements456.htm
- Gebhard, F. and Smalla, K., 1999. Monitoring field releases of genetically modified sugar beets for persistence of transgenic plant DNA and horizontal gene transfer. *FEMS Microbiology Ecology*, 28, 261-272.
- Goldstein, D.A., Tinland, B., Gilbertson, L.A., Staub, J.M., Bannon, G.A., Goodman, R.E., McCoy, R.L. and Silvanovich, A., 2005. Human safety and genetically modified plants: a review of antibiotic resistance markers and future transformation selection technologies. *Journal of Applied Microbiology*, 99, 7-23.
- Grant, R.J., Fanning, K.C., Kleinschmit, D., Stanisiewski, E.P., Hartnell, G.F., 2003. Influence of glyphosate-tolerant (event nk603) and corn rootworm protected (event MON863) corn silage and grain on feed consumption and milk production in Holstein cattle. *Journal of Dairy Science*, 86, 1707-15.
- Hammond, B., Lemen, J., Dudek, R., Ward, D., Jiang, C., Nemeth, M., Burns, J., 2006. Results of a 90-day safety assurance study with rats fed grain from corn rootworm-protected corn. *Food and Chemical Toxicology*, 44, 147-160.
- Head, G., Surber, J.B., Watson, J.A., Martin, J.W., Duan, J.J., 2002. No detection of Cry1Ac protein in soil after multiple years of transgenic Bt cotton (Bollguard) use. *Environmental Entomology*, 31, 30-36.
- Herman, R.A., S.L. Evans, D.M. Shanahan, C.A. Mihaliak, G.A. Bormett, D.L. Yound, and J. Buehrer, 2001. Rapid degradation of Cry1F delta-endotoxin in soil. *Environmental Entomology*, 30, 642-644.
- Herman, R.A., Wolt, J.D., Halliday, W.R., 2002. Rapid degradation of the Cry1F insecticidal crystal protein in soil. *Journal of Agricultural and Food Chemistry*, 50, 7076-7078.
- Hopkins, D.W., Gregorich, E.G., 2005. Decomposition of residues and loss of the δ -endotoxin from transgenic (Bt) corn (*Zea mays* L.) in soil. *Canadian Journal of Soil Science*, 85, 19-26.
- Icoz, I., Stotzky, G., 2007. Fate and effects of insect-resistant Bt crops in soil ecosystems. *Soil Biology and Biochemistry*, 40, 559-586.

- Kim, M.C., Ahn, J.H., Shin, H.C., Kim, T., Ryu, T.H., Kim, D.H., Song, H.G., Lee, G.H. and Ka, J.O., 2008. Molecular analysis of bacterial community structures in paddy soils for environmental risk assessment with two varieties of genetically modified rice, Iksan 483 and Milyang 204. *Journal of Microbiology and Biotechnology*, 18, 207-218.
- Krogh, H., Griffiths B., 2007. ECOGEN – Soil ecological and economic evaluation of genetically modified crops. *Pedobiologia*, 51, 171-173.
- Lawhorn, C.N., Neher, D.A., Dively, G.P., 2009. Impact of coleopteran targeting toxin (Cry3Bb1) of Bt corn on microbially mediated decomposition. *Applied Soil Ecology*, pre-print published online, DOI:10.1016/j.apsoil.2008.12.003.
- Lutz, B., Wiedemann, S., Einspanier, R., Mayer, J., Albrecht, C., 2005. Degradation of Cry1Ab Protein from Genetically Modified Maize in the Bovine Gastrointestinal Tract. *Journal of Agricultural and Food Chemistry*, 53, 1453-1456.
- Martinez, J.L., 2008. Antibiotics and antibiotic resistance genes in natural environments. *Science*, 321, 365-367.
- OECD, 1998. OECD Guidelines for the Testing of Chemicals / Section 4: Health Effects, Test No. 408: Repeated Dose 90-Day Oral Toxicity Study in Rodents. Organization for Economic Cooperation and Development, Paris.
- OIE, 2007. List of Antimicrobials of Veterinary importance. OIE International Committee. 75th general session, May 2007. Resolution no. XXVIII.
- Paget, E., Lebrun, M., Freyssinet, G. and Simonet, P., 1998. Fate of recombinant DNA in soil. *European Journal of Soil Biology*, 34, 81-88.
- Riesenfeld, C.S., Goodman, R.M. and Handelsman, J., 2004. Uncultured soil bacteria are a reservoir of new antibiotic resistance genes. *Environmental Microbiology*, 6, 981-989.
- Taylor, M.L., Hyun, Y., Hartnell, G.F., Riordan, S.G., Nemeth, M.A., Karunanandaa, K., George, B., Astwood, J.D., 2003. Comparison of broiler performance when fed diets containing grain from YieldGard Rootworm (MON863), YieldGard Plus (MON810 x MON863), nontransgenic control, or commercial reference corn hybrids. *Poultry Science*, 82, 1948-1956.
- WHO, 2007. Critically important antimicrobials for human medicine: Categorization for the development of risk management strategies to contain antimicrobial resistance due to non-human antimicrobial use. Report of the second WHO expert meeting, Copenhagen, 29-31 May 2007.
- WHO, 2005. Critically Important Antibacterial Agents for Human Medicine for Risk Management Strategies of Non-Human Use. Report of a WHO working group consultation, Canberra, 15-18 February 2005.
- Wiedemann, S., Lutz, B., Kurtz, H., Schwarz F. J., Albrecht, C., 2006. In situ studies on the time-dependent degradation of recombinant corn DNA and protein in the bovine rumen. *Journal of Animal Science*, 84, 135-144.

Wright, J.D., 2007. The antibiotic resistome: the nexus of chemical and genetic diversity.
Nature Reviews Microbiology, 5, 175-186.

SCIENTIFIC OPINION

Request from the European Commission related to the safeguard clause invoked by Austria on oilseed rape MS8, RF3 and MS8xRF3 according to Article 23 of Directive 2001/18/EC¹

Scientific Opinion of the Panel on Genetically Modified Organisms

(Question No EFSA-Q-2008-743)

Adopted on 15 June 2009

PANEL MEMBERS*

Hans Christer Andersson, Salvatore Arpaia, Detlef Bartsch, Josep Casacuberta, Howard Davies, Patrick du Jardin, Lieve Herman, Niels Hendriksen, Sirpa Kärenlampi, Jozsef Kiss, Gijs Kleter, Ilona Kryspin-Sørensen, Harry Kuiper, Ingolf Nes, Nickolas Panopoulos, Joe Perry, Annette Pötting, Joachim Schiemann, Willem Seinen, Jeremy Sweet and Jean-Michel Wal.

SUMMARY

On 15 July 2008, Austria invoked Article 23 of Directive 2001/18/EC (safeguard clause) to provisionally prohibit the marketing of genetically modified oilseed rape MS8, RF3 and MS8xRF3 on its territory. Austria provided detailed reasons listed in supporting documents.

On 31 July 2008, the European Food Safety Authority (EFSA) has been requested by the European Commission to provide a scientific opinion on the statement and documents submitted by Austria in the context of a safeguard clause invoked under Article 23 of Directive 2001/18/EC.

In light of the information package provided by Austria in support of its safeguard clause and, having considered all relevant scientific publications, the Scientific Panel on Genetically Modified Organisms (GMO Panel) of EFSA concludes that, in terms of risk to human and animal health and the environment, no new scientific evidence was presented that would invalidate the previous risk assessment of oilseed rape MS8, RF3 and MS8xRF3. The EFSA GMO Panel also concludes that no new scientific data or information was provided in support of adverse effects of oilseed rape MS8, RF3 and MS8xRF3 on the environment and on human

¹ For citation purposes: Scientific Opinion of the Panel on Genetically Modified Organisms on a request from the European Commission related to the safeguard clause invoked by Austria on oilseed rape MS8, RF3 and MS8xRF3 according to Article 23 of Directive 2001/18/EC. *The EFSA Journal* (2009) 1153, 1-16.

* (minority opinion) This opinion is not shared by 0 members of the Panel. / (conflict of interest) 0 members of the Panel did not participate in (part of) the discussion on the subject referred to above because of possible conflicts of interest.

and animal health in Austria. Therefore, no specific scientific evidence, in terms of risk to human and animal health and the environment, were provided that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC.

Key words: GMOs, oilseed rape (*Brassica napus*), MS8, RF3, MS8xRF3, Austria, safeguard clause, human and animal health, environment, Directive 2001/18/EC

TABLE OF CONTENTS

Panel Members	1
Summary	1
Table of Contents	3
Background	4
Terms of reference as provided by the European Commission.....	4
Acknowledgements	4
Assessment	5
1. Introduction	5
2. Assessment of documents provided by Austria.....	5
2.1. Food and feed safety issues.....	6
2.1.1. Toxicological and allergenicity assessment.....	6
2.2. Environmental safety issues and post-market environmental monitoring	6
2.2.1. Environmental risk assessment.....	6
2.2.2. Post-market environmental monitoring plan	9
2.2.3. Conclusion.....	9
Overall Conclusions and Recommendations.....	9
Documentation provided to EFSA	10
References	10

BACKGROUND

On 15 July 2008, Austria notified to the European Commission a national safeguard clause on genetically modified (GM) oilseed rape events MS8, RF3 and MS8xRF3 under Article 23 of Directive 2001/18/EC. The notification was accompanied by the scientific document entitled “*Scientific arguments for an import ban of herbicide tolerant oilseed rape Ms8, Rf3 and Ms8xRf3 (notification C/BE/96/01) in Austria*”.

On 31 July 2008, the European Food Safety Authority (EFSA) has received a request from the European Commission to provide a scientific opinion from its Scientific Panel on Genetically Modified Organisms (GMO Panel) on the statement and documents submitted by Austria in the context of its invoked safeguard clause.

TERMS OF REFERENCE AS PROVIDED BY THE EUROPEAN COMMISSION

EFSA was requested, under Article 29(1) and in accordance with Article 22(5) of Regulation (EC) No 178/2002, to provide a scientific opinion as to “*whether, in accordance with Article 23 of Directive 2001/18/EC, the statement and documents submitted by the Austrian authorities comprise new or additional information affecting the environmental risk assessment, such that detailed grounds exist to consider that the above authorised GMO, for the uses laid down in the corresponding consent, constitute a risk to human health or the environment*”.

ACKNOWLEDGEMENTS

The European Food Safety Authority wishes to thank the members of the Working Groups on Molecular Characterisation, Food/Feed and Environment, as well as the following members of its staff: Yann Devos, Yi Liu and Sylvie Mestdagh for the preparation of this opinion.

ASSESSMENT

1. Introduction

Directive 2001/18/EC provides the possibility for Member States to invoke safeguards on specific genetically modified organisms in the case where new or additional information, made available since the date of the consent, would affect the risk assessment of an authorised GMO. Provisions foreseen by Austria seek to provisionally prohibit the marketing of oilseed rape MS8, RF3 and MS8xRF3 for its intended uses on the Austrian territory.

The EFSA GMO Panel examined the set of supporting documents submitted by Austria. In this respect, the EFSA GMO Panel assessed whether the submitted documents comprise new scientific information that would change the outcome of previously performed risk assessments, and if detailed grounds exist to consider that the authorised oilseed rape MS8, RF3 and MS8xRF3, for its intended uses, constitute a risk to human and animal health or the environment.

The EFSA GMO Panel looked for evidence for GMO-specific risks – including long-term effects (e.g., BEETLE report, 2009) – taking into consideration the EFSA GMO Panel guidance document for the risk assessment of GM plants and derived food and feed (EFSA, 2006a) as well as any related risk assessments carried out in the past. In addition, the EFSA GMO Panel considered the relevance of raised concerns in light of the most recent scientific data and relevant peer-reviewed publications.

2. Assessment of documents provided by Austria

A set of supporting documents, accompanying the mandate of the European Commission (see *Terms of Reference as provided by the European Commission*), was forwarded to EFSA on 31 July 2008.

Austria provided the following set of documents in support of its safeguard clause:

- Austrian letter on new supplementary scientific evaluation (10 July 2008);
- Verbot des Inverkehrbringens von gentechnisch verändertem Raps aus den Ölrapslinien MS8, RF3 and MS8xRF3 in Österreich;
- Scientific arguments for an import ban of herbicide tolerant oilseed rape MS8, RF3 and MS8xRF3 (Notification C/BE/96/01) in Austria.

Based on the supporting documents, several issues were identified and therefore considered by the EFSA GMO Panel in the following two main areas: (1) the toxicological and allergenicity risk assessment, and (2) the environmental risk assessment and post-market environmental monitoring plan relating to the accidental spillage of oilseed rape MS8, RF3 and MS8xRF3 seeds.

Issues related to the coexistence of oilseed rape cropping systems and the adventitious presence of authorised GM material in non-GM products were not considered, as they fall outside the remit of the EFSA GMO Panel. In addition, the EFSA GMO Panel notes that it

only gives its opinion on the scientific quality of the post-market environmental monitoring activities proposed by applicants, whilst the final endorsement thereof is done by risk managers.

During its assessment, the EFSA GMO Panel identified issues raised by the Austrian authorities that would require further clarifications. To present and clarify the provided set of documents, an informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel and EFSA staff was held on 23 April 2009. A representative of the European Commission attended the meeting as observer.

2.1. Food and feed safety issues

2.1.1. Toxicological and allergenicity assessment

The EFSA GMO Panel observes that these two publications related to toxicological and allergenicity aspects of the risk assessment quoted by Austria (Spök et al., 2004, 2005) do not provide new data specific on the safety of oilseed rape MS8, RF3 and MS8xRF3: current approaches for assessing toxicological and allergenicity risks of genetically modified organisms are questioned in generic terms. The EFSA GMO Panel emphasises that the approach taken by the Panel in order to assess the potential toxicological and/or allergenicity risks of GM plants is in accordance with internationally developed guidelines (e.g., Codex Alimentarius, 2003).

Specific issues raised by Austria pertain to what they perceive as insufficient, inappropriate, or lacking data in the dossier. Moreover, Austria points to the observed differences in glucosinolate levels. The EFSA GMO Panel notes, however, that it has already taken the issues raised by Austria into account in its opinion on MS8xRF3, including the acute toxicity of phosphinothricin acetyltransferase (PAT), the safety of PAT if it would sustain food /feed processing, and variability in glucosinolate levels. The rabbit and chicken studies mentioned and considered inappropriate by Austria have not been included into the safety assessment of MS8xRF3 as summarized in the EFSA GMO Panel's opinion. Again, no new data indicating potential adverse effects of MS8xRF3 have thus been provided by Austria, whilst the data provided do not lead the EFSA GMO Panel to diverge from its previous opinion on the potential toxicity and allergenicity of MS8xRF3 (EFSA, 2005).

2.2. Environmental safety issues and post-market environmental monitoring

2.2.1. Environmental risk assessment

In line with its previous scientific opinions on herbicide tolerant oilseed rape GT73 (EFSA, 2004), MS8xRF3 (EFSA, 2005) and T45 (EFSA, 2008), the EFSA GMO Panel confirms that in regions where oilseed rape is grown and/or where oilseed rape seeds are imported and transported, feral oilseed rape populations are likely to occur in non-natural disturbed ecosystems (such as ports, processing facilities, margins of agricultural fields, roadside verges, railway lines, and wastelands) (Bagavathiannan and Van Acker, 2008). It is well-known that human activity contributes to the dispersal of plants (Wichmann et al., 2009), especially the transport of seeds by vehicles (Zwaenepoel et al., 2006; von der Lippe and Kowarik, 2007a,b; Garnier et al., 2008).

In the scientific literature, the occurrence of feral oilseed rape populations has been reported not only in Austria (Pascher et al., 2006), but also in Canada (Simard et al., 2002; Yoshimura et al., 2006; Knispel et al., 2008), France (Pessel et al., 2001; Garnier et al., 2008; Pivard et al., 2008a,b), Germany (Menzel, 2006; Reuter et al., 2008; Neuffer, 2009), Japan (Saji et al., 2005; Aono et al., 2006; Kawata et al., 2008; Nishizawa et al., 2009) and the United Kingdom (Crawley and Brown, 1995, 2004; Wilkinson et al., 1995; Charters et al., 1999; Norris and Sweet, 2002). These populations can be large and show significant variation in size from one year to the next (Crawley and Brown, 1995).

Due to its early germination potential and capacity to capture resources rapidly, oilseed rape can take advantage of disturbed land (Blackshaw et al., 2003, 2004). However, successful recruitment of oilseed rape from seed spillage from passing traffic mainly depends upon its ability to compete for space with primary colonizers, particularly perennial grasses. In most non-agricultural areas, oilseed rape lacks the ability to establish stable populations due to the absence of competition-free gaps (Crawley et al., 1993; Warwick et al., 1999; Hails et al., 2006). Once established, oilseed rape populations often become extinct after 2 to 4 years (Crawley and Brown, 1995; Crawley et al., 2001; Norris and Sweet, 2002). If habitats are disturbed on a regular basis (e.g., by mowing, herbicide application, soil disturbance) and replenished with seed from seed spillage or recruitment from seeds produced by residents or from seeds from the seedbank, then feral oilseed rape populations can persist for longer periods (8-10 years) (Pessel et al., 2001; Pivard et al., 2008a,b). Using genetic analyses and farmer surveys, Pessel et al. (2001) revealed that some members of feral oilseed rape populations in road verges in France originated from varieties that had not been marketed for at least 8 years.

Oilseed rape is generally regarded as an opportunistic species, and not as an environmentally hazardous colonizing species (Warwick et al., 1999). Several field studies and model predictions reported that the presence of herbicide tolerance in oilseed rape does not confer a fitness advantage, unless the respective herbicide is applied (Crawley et al., 1993, 2001; Fredshavn et al., 1995; Warwick et al., 1999, 2004; Norris and Sweet, 2002; Claessen et al., 2005a,b; Simard et al., 2005; Garnier and Lecomte, 2006; Garnier et al., 2006). In the absence of glufosinate-ammonium-containing herbicide applications, oilseed rape MS8, RF3 and MS8xRF3 is neither more likely to survive, nor more invasive or persistent than its conventional counterpart. Moreover, there is no evidence that tolerance to glufosinate-ammonium enhances seed dormancy, and hence the persistence of feral oilseed rape populations (Crawley et al., 1993, 2001; Claessen et al., 2005a,b; Sweet et al., 2004; Lutman et al., 2005; Messéan et al., 2007). Because glufosinate-ammonium-containing herbicides are not widely used in ruderal ecosystems in the European Union (EU), feral oilseed rape plants ensuing from spilled seeds of oilseed rape MS8, RF3 and MS8xRF3 would not show any enhanced fitness and would thus behave as conventional plants. Only where and when glufosinate-ammonium-containing herbicides are applied, is oilseed rape MS8, RF3 and MS8xRF3 expected to have a fitness advantage. Likewise, there are no indications that the *barstar/barnase* gene complex would alter seed survival characteristics and confer a selective advantage (Fredshavn et al., 1995; Sweet et al., 2004; Lutman et al., 2008). The scientific information provided in the Austrian safeguard clause notification does not give any new information regarding increased likelihood of establishment or survival of feral oilseed rape plants in case of accidental release into the environment of oilseed rape MS8, RF3 and MS8xRF3 seeds during transportation and processing.

The EFSA GMO Panel is aware that if feral oilseed rape plants derived from spilled seeds remain uncontrolled and reproduce, they may survive, outcross and eventually disperse genes to cross-compatible plants such as *Brassica rapa* and *Raphanus raphanistrum* (Scheffler and Dale, 1994; Eastham and Sweet, 2002; Chèvre et al., 2004; Warwick et al., 2003, 2004, 2008; Claessen et al., 2005b; Jørgensen, 2007; BEETLE report, 2009; Devos et al., 2009; Jørgensen et al., 2009). Scientific evidence suggests that feral oilseed rape populations may serve as a reservoir that could hold and return (trans)genes to cultivated populations of oilseed rape in a different place and time, and act as a genetic bridge delivering the (trans)genes to sympatric cross-compatible plants (Saji et al., 2005; Aono et al., 2006; Pascher et al., 2006; Yoshimura et al., 2006; Knispel et al., 2008; Nishizawa et al., 2009). In Canadian regions where GM oilseed rape is frequently grown (e.g., Beckie et al., 2006), feral oilseed rape populations were shown to actively outcross with cultivated populations of GM oilseed rape and to accumulate transgenes (Knispel et al., 2008). However, compared to cultivated oilseed rape populations, the contribution of feral oilseed rape plants in vertical gene flow is expected to be limited: feral oilseed rape populations are small compared to cultivated populations and contribute little to the pollen load in the environment (Colbach et al., 2001a,b, 2005; Devaux et al., 2005, 2007, 2008; Gruber and Claupein, 2006; Knispel et al., 2008; Colbach, 2009). Moreover, there are no compelling data to suggest that the presence of an herbicide tolerance trait in a wild relative changes the behaviour of the wild relative so far (e.g., Warwick et al., 2008). In the absence of glufosinate-ammonium-containing herbicides, hybrids or wild relatives containing the herbicide tolerance trait do not show any enhanced fitness and behave as conventional plants. Thus escaped plants and genes dispersed to other cross-compatible plants would not create additional environmental impacts. If needed, feral oilseed rape MS8xRF3 and hybridised/introgressed relatives can be managed by the use of other herbicides and/or adequate mechanical practices (Beckie et al., 2004; Devos et al., 2004; Warwick et al., 2004; Légère, 2005; Simard et al., 2005; Gruber et al., 2008; Lutman et al., 2008).

The environmental exposure due to GM oilseed rape grain imports is anticipated to be low, as the amounts of viable oilseed rape seeds imported in the EU are limited with most seeds being imported by boat and crushed in or near the ports of entry. Some of the oilseed rape seeds imported into the EU are likely to be transported inland to Austria by boat. Moreover, some oilseed rape seeds entering Austria are transported by road or rail to processing plants in Austria. As indicated above, survival and outcrossing from plants derived from seed spillage will be at very low frequencies and have no hazardous environmental consequences compared to current feral oilseed rape populations.

In conclusion, the EFSA GMO Panel confirms that feral oilseed rape plants are likely to occur wherever oilseed rape is cultivated and/or transported and that transgenic oilseed rape is no exception (e.g., Saji et al., 2005; Aono et al., 2006; Yoshimura et al., 2006; Bagavathiannan and Van Acker, 2008; Kawata et al., 2008; Knispel et al., 2008; Nishizawa et al., 2009). However, there is no evidence that the herbicide tolerance trait introduced by genetic engineering results in increased invasiveness of oilseed rape MS8, RF3 and MS8xRF3, except when glufosinate-ammonium-containing herbicides are applied. As such, escaped plants and genes dispersed to other cross-compatible plants would not create additional agronomic or environmental impacts. This – together with the assessment that oilseed rape MS8, RF3 and MS8xRF3 and hybridising relatives have no enhanced fitness or invasiveness characteristics (except in the presence of glufosinate-ammonium-containing herbicides) – confirms earlier conclusions of the EFSA GMO Panel.

2.2.2. Post-market environmental monitoring plan

The EFSA GMO Panel maintains its position that the scope of the post-market environmental monitoring plan provided by the applicant complies with (1) the intended uses of oilseed rape MS8, RF3 and MS8xRF3, which excludes cultivation, (2) the requirements of the EFSA GMO Panel guidance document for the risk assessment of GM plants and food and feed products (EFSA, 2006a), and (3) the EFSA GMO Panel scientific opinion on post-market environmental monitoring (EFSA, 2006b).

European operators importing, handling and processing viable oilseed rape commodities have recently joined with the European Association of Bioindustries (EuropaBio) in developing monitoring systems for GM oilseed rape imported through main points of entry and processing facilities (Lecoq et al., 2007; Windels et al., 2008). These monitoring systems are based on Hazard Analysis and Critical Control Points (HACCP) principles. Therefore, it is anticipated that risk managers will opt for post-market environmental monitoring of imported oilseed rape MS8, RF3 and MS8xRF3 seeds in accordance with these arrangements.

2.2.3. Conclusion

The EFSA GMO Panel confirms its opinion that the likelihood of unintended environmental effects as a consequence of spread of (trans)genes from oilseed rape MS8, RF3 and MS8xRF3 will not differ from that of conventional oilseed rape varieties in the context of its intended uses. The EFSA GMO Panel agrees with the monitoring plan submitted by the applicant, especially now that comprehensive arrangements have been made by applicants and operators for monitoring at major points of import and processing in the EU. However, the EFSA GMO Panel continues to recommend that appropriate management systems should be in place to minimise accidental loss and spillage of transgenic oilseed rape seeds during transportation, storage and handling in the environment, and processing into derived products. These conclusions are in line with previous scientific opinions of the EFSA GMO Panel on the import and processing of herbicide-tolerant oilseed rape GT73 (EFSA, 2004), MS8xRF3 (EFSA, 2005) and T45 (EFSA, 2008).

OVERALL CONCLUSIONS AND RECOMMENDATIONS

The EFSA GMO Panel has investigated the claims and documents submitted in support of the Austrian safeguard clause and presented at the informal meeting between the Austrian delegation, several experts of the EFSA GMO Panel and EFSA staff on 23 April 2009. In these documents, the EFSA GMO Panel did not identify any new data subject to scientific scrutiny or scientific information that would change previous risk assessments conducted on oilseed rape MS8, RF3 and MS8xRF3 which currently has marketing consent in the EU. Furthermore, the Austrian submission did not supply scientific evidence, that the environment or ecology of Austria was different from other regions of the EU, sufficient to merit separate risk assessments from those conducted for other regions in the EU.

Having considered the overall information package submitted by Austria as well as a broad range of relevant scientific literature, the EFSA GMO Panel is of the opinion that there is no specific scientific evidence, in terms of risk to human and animal health and the environment, that would justify the invocation of a safeguard clause under Article 23 of Directive 2001/18/EC for the marketing of oilseed rape MS8, RF3 and MS8xRF3 for its intended uses in Austria. In conclusion, the EFSA GMO Panel finds that the scientific evidence currently

available does not sustain the arguments provided by Austria, and therefore the EFSA GMO Panel reiterates its previous scientific opinion on oilseed rape MS8, RF3 and MS8xRF3.

DOCUMENTATION PROVIDED TO EFSA

1. Letter, dated 31 July 2008, with supporting documents from Jos Delbeke, Acting Director-General Environment EC, to Catherine Geslain-Lanéelle, Executive Director EFSA (ref ENV/B3/AA/lh ARES(2008) 18278), requesting for a scientific opinion on the safeguard notification submitted by Austria under Article 23 of Directive 2001/18/EC for oilseed rape MS8, RF3 and MS8xRF3 and comprising the following supporting documents:
 - Scientific arguments for an import ban of herbicide tolerant oilseed rape MS8, RF3 and MS8xRF3 (notification C/BE/96/01) in Austria.
2. Letter, dated 15 October 2008, from Catherine Geslain-Lanéelle, Executive Director EFSA to Jos Delbeke, Director-General Environment EC (ref CGL/RM/PB/SM/shv (2008) 3369831), acknowledging the receipt of the mandate accompanied with the supporting documents.

REFERENCES

- Aono, M., Wakiyama, S., Nagatsu, M., Nakajima, N., Tamaoki, M., Kubo, A., Saji, H., 2006. Detection of feral transgenic oilseed rape with multiple-herbicide resistance in Japan. *Environmental Biosafety Research*, 5: 77-87.
- Bagavathiannan, M.V., Van Acker, R.C., 2008. Crop ferality: implications for novel trait confinement. *Agriculture, Ecosystems and Environment*, 127: 1-6.
- Beckie, H.J., Séguin-Swartz, G., Nair, H., Warwick, S.I., Johnson, E., 2004. Multiple herbicide-resistant canola (*Brassica napus*) can be controlled by alternative herbicides. *Weed Science*, 52: 152-157.
- Beckie, H.J., Harker, K.N., Hall, L.M., Warwick, S.I., Légère, A., Sikkema, P.H., Clayton, G.W., Thomas, A.G., Leeson, J.Y., Séguin-Swartz, G., Simard, M.-J., 2006. A decade of herbicide-resistant crops in Canada. *Canadian Journal of Plant Science*, 86: 1243-1264.
- BEETLE report, 2009. Long-term effects of genetically modified (GM) crops on health and the environment (including biodiversity): prioritization of potential risks and delimitation of uncertainties. German Federal Office of Consumer Protection and Food Safety, BLaU-Umweltstudien and Genius GmbH, http://ec.europa.eu/environment/biotechnology/pdf/beetle_report.pdf
- Blackshaw, R.E., Brandt, R.N., Janzen, H.H., Entz, T., Grant, C.A., Derksen, D.A., 2003. Differential response of weed species to added nitrogen. *Weed Science*, 51: 532-539.
- Blackshaw, R.E., Brandt, R.N., Janzen, H.H., Entz, T., 2004. Weed species response to phosphorus fertilization. *Weed Science*, 52: 406-412.
- Charters, Y.M., Robertson, A., Squire, G.R., 1999. Investigation of feral oilseed rape populations, genetically modified organisms research report (No. 12), Department of the

Environment, Transport and the Regions,
<http://www.defra.gov.uk/environment/gm/research/reports.htm>

- Chèvre, A.M., Ammitzbøll, H., Breckling, B., Dietz-Pfeilstetter, A., Eber, F., Fargue, A., Gomez-Campo, C., Jenczewski, E., Jørgensen, R., Lavigne, C., Meier, M., den Nijs, H., Pascher, K., Seguin-Swartz, G., Sweet, J., Stewart, N., Warwick, S., 2004. A review on interspecific gene flow from oilseed rape to wild relatives. In: den Nijs, H.C.M., Bartsch, D., Sweet, J. (Eds), *Introgression from Genetically Modified Plants into Wild Relatives* (pp 235-251), CABI publishing.
- Claessen, D., Gilligan, C.A., Lutman, P.J.W., van den Bosch, F., 2005a. Which traits promote persistence of feral GM crops? Part 1: implications of environmental stochasticity. *Oikos*, 110: 20-29.
- Claessen, D., Gilligan, C.A., van den Bosch, F., 2005b. Which traits promote persistence of feral GM crops? Part 2: implications of metapopulation structure. *Oikos*, 110: 30-42.
- Codex Alimentarius, 2003. Codex principles and guidelines on foods derived from biotechnology. Joint FAO/WHO Food Standards Programme, Food and Agriculture Organisation, Rome.
- Colbach, H., 2009. How to model and simulate the effects of cropping systems on population dynamics and gene flow at the landscape level: example of oilseed rape volunteers and their role for co-existence of GM and non-GM crops. *Environmental Science and Pollution Research*, 16: 348-360.
- Colbach, N., Clermont-Dauphin, C., Meynard, J.M., 2001a. GeneSys: a model of the influence of cropping system on gene escape from herbicide tolerant rapeseed crops to rape volunteers I. Temporal evolution of a population of rapeseed volunteers in a field. *Agriculture, Ecosystems and Environment*, 83: 235-254.
- Colbach, N., Clermont-Dauphin, C., Meynard, J.M., 2001b. GeneSys: a model of the influence of cropping system on gene escape from herbicide tolerant rapeseed crops to rape volunteers II. Genetic exchanges among volunteer and cropped population a small region. *Agriculture, Ecosystems and Environment*, 83: 255-270.
- Colbach, N., Molinari, N., Meynard, J.M., Messéan, A., 2005. Spatial aspects of gene flow between rapeseed varieties and volunteers. *Agronomy for Sustainable Development*, 25: 355-368.
- Crawley, M.J., Brown, S.L., 1995. Seed limitation and the dynamics of feral oilseed rape on the M25 motorway. *Proceedings of the Royal Society B-Biological Sciences*, 259: 49-54.
- Crawley, M.J., Brown, S.L., 2004. Spatially structured population dynamics in feral oilseed rape. *Proceedings of the Royal Society B-Biological Sciences*, 271: 1909-1916.
- Crawley, M.J., Hails, R.S., Rees, M., Kohn, D., Buxton, J., 1993. Ecology of transgenic oilseed rape in natural habitats. *Nature*, 363: 620-623.
- Crawley, M.J., Brown, S.L., Hails, R.S., Kohn, D.D., Rees, M., 2001. Transgenic crops in natural habitats. *Nature*, 409: 682-683.

- Devaux, C., Lavigne, C., Falentin-Guyomarc'h, H., Vautrin, S., Lecomte, J., Klein, E.K., 2005. High diversity of oilseed rape pollen clouds over an agro-ecosystem indicated long-distance dispersal. *Molecular Ecology*, 14: 2269-2280.
- Devaux, C., Lavigne, C., Austerlitz, F., Klein, E.K., 2007. Modelling and estimating pollen movement in oilseed rape (*Brassica napus*) at the landscape scale using genetic markers. *Molecular Ecology*, 16: 487-499.
- Devaux, C., Klein, E.K., Lavigne, C., Sausse, C., Messéan, A., 2008. Environmental and landscape effects on cross-pollination rates observed at the long distance among French oilseed rape (*Brassica napus*) commercial fields. *Journal of Applied Ecology*, 45: 803-812.
- Devos, Y., Reheul, D., De Schrijver, A., Cors, F., Moens, W., 2004. Management of herbicide-tolerant oilseed rape in Europe: a case study on minimizing vertical gene flow. *Environmental Biosafety Research*, 3: 135-148.
- Devos, Y., De Schrijver, A., Reheul, D., 2009. Quantifying the introgressive hybridisation propensity between transgenic oilseed rape and its wild/weedy relatives. *Environmental Monitoring and Assessment*, 149: 303-322.
- Eastham, K., Sweet, J., 2002. Genetically modified organisms (GMOs): the significance of gene flow through pollen transfer, European Environment Agency, http://www.eea.europa.eu/publications/environmental_issue_report_2002_28
- EFSA, 2004. Opinion of the Scientific Panel on Genetically Modified Organisms on a request from the Commission related to the Notification (Reference C/NL/98/11) for the placing on the market of herbicide-tolerant oilseed rape GT73, for import and processing, under Part C of Directive 2001/18/EC from Monsanto. *The EFSA Journal*, 29: 1-19, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620772413.htm
- EFSA, 2005. Opinion of the Scientific Panel on Genetically Modified Organisms on a request from the Commission related to the application (Reference C/BE/96/01) for the placing on the market of glufosinate-tolerant hybrid oilseed rape Ms8 x Rf3, derived from genetically modified parental lines (Ms8, Rf3), for import and processing for feed and industrial uses, under Part C of Directive 2001/18/EC from Bayer CropScience. *The EFSA Journal*, 281: 1-23, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620770114.htm
- EFSA, 2006a. Guidance document of the Scientific Panel on Genetically Modified Organisms for the risk assessment of genetically modified plants and derived food and feed. *The EFSA Journal*, 99: 1-100, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620775747.htm
- EFSA, 2006b. Opinion of the Scientific Panel on Genetically Modified Organisms on the Post Market Environmental Monitoring (PMEM) of genetically modified plants. *The EFSA Journal*, 319: 1-27, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620769727.htm
- EFSA, 2008. Opinion of the Scientific Panel on Genetically Modified Organisms on an application (Reference EFSA-GMO-UK-2005-25) for the placing on the market of

glufosinate-tolerant oilseed rape T45 for food and feed uses, import and processing and renewal of the authorization of oilseed rape T45 as existing products, both under Regulation (EC) 1829/2003 from Bayer CropScience. *The EFSA Journal*, 635: 1-22, http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178690393760.htm

- Fredshavn, J.R., Poulsen, G., Huybrechts, I., Rüdelsheim, P., 1995. Competitiveness of transgenic oilseed rape. *Transgenic Research*, 4: 142-148.
- Garnier, A., Lecomte, J., 2006. Using spatial and stage-structured invasion model to assess the spread of feral population of transgenic oilseed rape. *Ecological Modelling*, 194: 141-149.
- Garnier, A., Deville, A., Lecomte, J., 2006. Stochastic modelling of feral plant populations with seed immigration and road verge management. *Ecological Modelling*, 197: 373-382.
- Garnier, A., Pivard, S., Lecomte, J., 2008. Measuring and modelling anthropogenic secondary seed dispersal along roadverges for feral oilseed rape. *Basic and Applied Ecology*, 9: 533-541.
- Gruber, S., Claupein, W., 2006. Fecundity of volunteer oilseed rape and estimation of potential gene dispersal by a practice-related model. *Agriculture, Ecosystems and Environment*, 119: 401-408.
- Gruber, S., Colbach, N., Barbottin, A., Pekrun, C., 2008. Post-harvest gene escape and approaches for minimizing it. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*, 3: 1-17.
- Hails, R.S., Bullock, J.M., Morley, K., Lamb, C., Bell, P., Horsnell, R., Hodgson, D.J., Thomas, J., 2006. Predicting fitness changes in transgenic plants: testing a novel approach with pathogen resistant Brassicas. *IOBC/WPRS Bulletin*, 29: 63-70.
- Jørgensen, R.B., 2007. Oilseed rape: Co-existence and gene flow from wild species. *Advances in Botanical Research*, 45: 451-464.
- Jørgensen, R.B., Hauser, T., D'Hertefeldt, T., Andersen, N.S., Hooftman, D., 2009. The variability of processes involved in transgene dispersal – case studies from *Brassica* and related genera. *Environmental Science and Pollution Research*, pre-print published online, DOI:10.1007/s11356-009-0142-4.
- Kawata, M., Murakami, K., Ishikawa, T., 2008. Dispersal and persistence of genetically modified oilseed rape around Japanese harbors. *Environmental Science and Pollution Research*, 16: 120-126.
- Knispel, A.L., McLachlan, S.M., Van Acker, R.C., Friesen, L.F., 2008. Gene flow and multiple herbicide resistance in escaped canola populations. *Weed Science*, 56: 72-80.
- Lecoq, E., Holt, K., Janssens, J., Legris, G., Pleysier, A., Tinland, B., Wandelt, C., 2007. General surveillance: roles and responsibilities the industry view. *Journal of Consumer Protection and Food Safety*, 2(S1): 25-28.

- Légère, A., 2005. Risks and consequences of gene flow from herbicide-resistant crops: canola (*Brassica napus* L.) as a case study. *Pest Management Science*, 61: 292-300.
- Lutman, P.J.W., Berry, K., Payne, R.W., Simpson, E., Sweet, J.B., Champion, G.T., May, M.J., Wightman, P., Walker, K., Lainsbury, M., 2005. Persistence of seeds from crops of conventional and herbicide tolerant oilseed rape (*Brassica napus*). *Proceedings of the Royal Society B-Biological Sciences*, 272: 1909-1915.
- Lutman, P.J.W., Sweet, J., Berry, K., Law, J., Payne, R., Simpson, E., Walker, K., Wightman, P., 2008. Weed control in conventional and herbicide tolerant winter oilseed rape (*Brassica napus*) grown in rotations with winter cereals in the UK. *Weed Research*, 48: 408-419.
- Menzel, G., 2006. Verbreitungsdynamik und Auskreuzungspotential von *Brassica napus* L. (Raps) im Großraum Bremen. GCA-Verlag, Waabs, ISBN 3-89863-213-X.
- Messéan, A., Sausse, C., Gasquez, J., Darmency, H., 2007. Occurrence of genetically modified oilseed rape seeds in the harvest of subsequent conventional oilseed rape over time. *European Journal of Agronomy*, 27: 115-122.
- Neuffer, B., 2009. Kulturpflanzen als Quelle für Neophyten – Etablierungsmechanismen bei der Verwilderung von Raps (2004-2006), und Abschätzung des Hybridisierungspotentials ruderaler Raps- und Rübsenpopulationen (2007-2009) Im Rahmen des BIOLOG EUROPA Projektverbundes "Evolutionäre, ökologische und gesellschaftliche Konsequenzen biologischer Invasionen", http://www.gmo-safety.eu/en/safety_science/188.docu.html
- Nishizawa, T., Nakajima, N., Aono, M., Tamaoki, M., Kubo, A., Saji, H., 2009. Monitoring the occurrence of genetically modified oilseed rape growing along a Japanese roadside: 3-year observations. *Environmental Biosafety Research*, 8: 34-44.
- Norris, C., Sweet, J., 2002. Monitoring large scale releases of genetically modified crops (EPG1/5/84) incorporating report on project EPG 1/5/30: monitoring releases of genetically modified crop plants, DEFRA report, EPG 1/5/84, http://www.defra.gov.uk/environment/gm/research/pdf/epg_1-5-84_screen.pdf
- Pascher, K., Narendja, F., Rau, D., 2006. Feral oilseed rape – Investigations on its potential for hybridisation, Studie im Auftrag des Bundesministeriums fuer Gesundheit und Frauen, Forschungsberichte der Sektion IV, Band 3/2006, http://www.bmgfj.gv.at/cms/site/attachments/8/1/9/CH0255/CMS1138950978238/feral_oilseed_rape_-_investigation_on_its_potential_for_hybridisation_gesamt_f_hp.pdf
- Pessel, F.D., Lecomte, J., Emeriau, V., Krouti, M., Messéan, A., Gouyon, P.H., 2001. Persistence of oilseed rape (*Brassica napus* L.) outside of cultivated fields. *Theoretical and Applied Genetics*, 102: 841-846.
- Pivard, S., Adamczyk, K., Lecomte, J., Lavigne, C., Bouvier, A., Deville, A., Gouyon, P.H., Huet, S., 2008a. Where do the feral oilseed rape populations come from? A large-scale study of their possible origin in a farmland area. *Journal of Applied Ecology*, 45: 476-485.

- Pivard, S., Demšar, D., Lecomte, J., Debeljak, M., Džeroski, S., 2008b. Characterizing the presence of oilseed rape feral populations on field margins using machine learning. *Ecological Modelling*, 212: 147-154.
- Reuter, H., Menzel, G., Pehlke, H., Breckling, B., 2008. Hazard mitigation or mitigation hazard? Would genetically modified dwarfed oilseed rape (*Brassica napus*) increase feral survival? *Environmental Science and Pollution Research*, 15: 529-535.
- Saji, H., Nakajima, N., Aono, M., Tamaoki, M., Kubo, A., Wakiyama, S., Hatase, Y., Nagatsu, M., 2005. Monitoring the escape of transgenic oilseed rape around Japanese ports and roadsides. *Environmental Biosafety Research*, 4: 217-222.
- Scheffler, J.A., Dale, P.J., 1994. Opportunities for gene transfer from transgenic oilseed rape (*Brassica napus*) to related species. *Transgenic Research*, 3: 263-278.
- Simard, M.J., Légère, A., Pageau, D., Lajeunesse, J., Warwick, S., 2002. The frequency and persistence of canola (*Brassica napus*) volunteers in Québec cropping systems. *Weed Technology*, 16: 433-439.
- Simard, M.J., Légère, A., Séguin-Swartz, G., Nair, H., Warwick, S., 2005. Fitness of double vs. single herbicide-resistant canola. *Weed Science*, 53: 489-498.
- Spök, A., Hofer, H., Lehner, P., Valenta, R., Stirn, S., Gaugitsch, H., 2004. Risk assessment of GMO products in the European Union. Umweltbundesamt Wien. Berichte, Band 253. July 2004, http://www.umweltbundesamt.at/en/publikationen/publikationssuche/publikationsdetail/?pub_id=1531
- Spök, A., Gaugitsch, H., Laffer, S., Pauli, G., Saito, H., Sampson, H., Sibanda, E., Thomas, W., Van Hage, M., Valenta, R., 2005. Suggestions for the assessment of the allergenic potential of genetically modified organisms. *International Archives of Allergy and Immunology*, 137: 167-180.
- Sweet, J., Simpson, E., Law, J., Lutma, P., Berry, K., Payne, R., Champion, G., May, M., Walker, K., Wightman, P., Lainsbury, M., 2004. Botanical and Rotational Implications of Genetically Modified Herbicide Tolerance (BRIGHT) HGCA Project Report 353, 265. HGCA London, UK.
- von der Lippe, M., Kowarik, I., 2007a. Long-distance dispersal of plants by vehicles as a driver of plant invasions. *Conservation Biology*, 21: 986-996.
- von der Lippe, M., Kowarik, I., 2007b. Crop seed spillage along roads: a factor of uncertainty in the containment of GMO. *Ecography*, 30: 483-490.
- Warwick, S.I., Beckie, H.J., Small, E., 1999. Transgenic crops: new weed problems for Canada? *Phytoprotection*, 80: 71-84.
- Warwick, S., Beckie, H.J., Simard, M.J., Légère, A., Nair, H., Séguin-Swartz, G., 2004. Environmental and agronomic consequences of herbicide-resistant (HR) canola in Canada. In: den Nijs, H.C.M., Bartsch, D., Sweet, J. (Eds), *Introgression from Genetically Modified Plants into Wild Relatives* (pp 323-337), CABI publishing.

- Warwick, S.I., Simard, M.J., Légère, A., Beckie, H.J., Braun, L., Zhu, B., Mason, P., Séguin-Swartz, G., Stewart, C.N. Jr., 2003. Hybridization between transgenic *Brassica napus* L. and its wild relatives: *B. rapa* L., *Raphanus raphanistrum* L., *Sinapis arvensis* L., and *Erucastrum gallicum* (Willd.) O.E. Schulz. *Theoretical and Applied Genetics*, 107: 528-539.
- Warwick, S.I., Légère, A., Simard, M.-J., James, T., 2008. Do escaped transgenes persist in nature? The case of an herbicide resistance transgene in a weedy *Brassica rapa* population. *Molecular Ecology*, 17: 1387-1395.
- Wichmann, M.C., Alexander, M.J., Soons, M.B., Galsworthy, S., Dunne, L., Gould, R., Fairfax, C., Niggemann, M., Hails, R.S., Bullock, J.M., 2009. Human-mediated dispersal of seeds over long distances. *Proceedings of the Royal Society B-Biological Sciences*, 276: 523-532.
- Wilkinson, M.J., Timmons, A.M., Charters, Y., Dubbels, S., Robertson, A., Wilson, N., Scott, S., O'Brien, E., Lawson, H.M., 1995. Problems of risk assessment with genetically modified oilseed rape. In: *Proceedings of the Brighton Crop Protection Conference – Weeds*, 3: 1035-1044.
- Windels, P., Alcalde, E., Lecoq, E., Legris, G., Pleysier, A., Tinland, B., Wandelt, C., 2008. General surveillance for import and processing: the EuropaBio approach. *Journal of Consumer Protection and Food Safety*, 3(S2): 14-16.
- Yoshimura, Y., Beckie, H.J., Matsuo, K., 2006. Transgenic oilseed rape along transportation routes and port of Vancouver in western Canada. *Environmental Biosafety Research*, 5: 67-75.
- Zwaenepoel, A., Roovers, P., Hermy, M., 2006. Motor vehicles as vectors of plant species from road verges in a suburban environment. *Basic and Applied Ecology*, 7: 83-93.